

## CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation of U.S. Application No. 10/411, 484, filed April 8, 2003 which is a continuation-in-part of U.S. Application No. 10/302,124, filed November 22, 2002, which is a continuation-in-part of U.S. Application No. 10/266,213 filed October 8, 2002, which is a continuation of U.S. 09/610,456, filed July 5, 2000, now U.S. 6,472,406, which claims the benefit of U.S. Provisional Application No. 60/142,362, filed on July 6, 1999.

## BACKGROUND OF THE INVENTION

Field of the Invention

[0002] The invention relates to bacterial antibiotic resistance. More particularly, the invention relates to compositions and methods for overcoming bacterial antibiotic resistance

Brief Summary of the Related Art

[0003] Bacterial antibiotic resistance has become one of the most important threats to modern health care. Cohen, Science 257:1051-1055 (1992) discloses that infections caused by resistant bacteria frequently result in longer hospital stays, higher mortality and increased cost of treatment. Neu, Science 257:1064-1073 (1992) discloses that the need for new antibiotics will continue to escalate because bacteria have a remarkable ability to develop resistance to new agents rendering them quickly ineffective.

[0004] The present crisis has prompted various efforts to elucidate the mechanisms responsible for bacterial resistance. Coulton *et al.*, Progress in Medicinal Chemistry 31:297-349 (1994) teaches that the widespread use of penicillins and cephalosporins has resulted in the emergence of  $\beta$ -lactamases, a family of bacterial enzymes that catalyze the hydrolysis of the  $\beta$ -lactam ring common to numerous presently used antibiotics. More recently, Dudley, Pharmacotherapy 15: 9S-14S (1995) has disclosed that resistance mediated by  $\beta$ -lactamases is a critical aspect at the core of the development of bacterial antibiotic resistance.

[0005] Attempts to address this problem through the development of  $\beta$ -lactamase inhibitors have had limited success. Sutherland, Trends Pharmacol. Sci. 12: 227-232 (1991) discusses the development of the first clinically useful  $\beta$ -lactamase inhibitor, clavulanic acid, which is a metabolite of *Streptomyces clavuligerus*. Coulton *et al* (supra) discloses two semi-synthetic inhibitors, sulbactam and tazobactam, presently available. Coulton *et al.* (supra) also teaches that in combination with  $\beta$ -lactamase-susceptible antibiotics,  $\beta$ -lactamase inhibitors prevent antibiotic inactivation by  $\beta$ -lactamase enzymes, thereby producing a synergistic effect against  $\beta$ -lactamase producing bacteria

[0006] Li et al., Bioorg. Med. Chem. 5 (9): 1783-1788 (1997), discloses that  $\beta$ -lactamase enzymes are inhibited by phosphonate monoesters. Li et al. teaches that better inhibitory activity is achieved by compounds with amido side-chains, but that such compounds suffer the disadvantage of hydrolytic instability. Li et al. discloses that benzyisulfonamidomethyl phosphonate monoesters exhibit better hydrolytic stability, but also significantly weaker potency against  $\beta$ -lactamase enzymes, than do the corresponding benzylamidomethyl phosphonate monoesters. Dryjanski and Pratt, Biochemistry 34:3569-3575 (1995) teaches that *p*-nitrophenyl [(dansylamido)methyl]phosphonate irreversibly inactivates the P99  $\beta$ -lactamase enzyme, and describes its use as a mechanistic probe for studying the interaction of ligands with a second binding site of the enzyme.

[0007] The availability of only a few  $\beta$ -lactamase inhibitors, however, is insufficient to counter the constantly increasing diversity of  $\beta$ -lactamases, for which a variety of novel and distinct inhibitors has become a necessity. There is, therefore, a need for the ability to identify new  $\beta$ -lactamase inhibitors. The development of fully synthetic inhibitors would greatly facilitate meeting this need.

### BRIEF SUMMARY OF THE INVENTION

[0008] The invention provides novel  $\beta$ -lactamase inhibitors, which are structurally unrelated to the natural product and semi-synthetic  $\beta$ -lactamase inhibitors presently available, and which do not require a  $\beta$ -lactam pharmacophore.

[0009] In a second aspect, the invention provides pharmaceutical compositions comprising a compound of the invention and a pharmaceutically acceptable carrier or diluent.

[0010] In a third aspect, the invention provides methods for inhibiting bacterial growth, such methods comprising administering to a bacterial cell culture, or to a bacterially infected cell culture, tissue, or organism, a  $\beta$ -lactamase inhibitor of the invention.

### BRIEF DESCRIPTION OF THE FIGURES

[0011] Figure 1 is a graphical plot of the synergistic effect of  $\beta$ -lactamase inhibitors as a function of log P. Synergy is defined and methods for its determination are provided in the Examples.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0012] The invention relates to bacterial antibiotic resistance. More particularly, the invention relates to compositions and methods for overcoming bacterial antibiotic resistance. The patents and publications identified in this specification indicate the knowledge in this field and are hereby incorporated by reference in their entirety. In the case of inconsistencies, the present disclosure will prevail.

**[0013]** The invention provides novel  $\beta$ -lactamase inhibitors, which are structurally unrelated to the natural product and semi-synthetic  $\beta$ -lactamase inhibitors presently available, and which do not require a  $\beta$ -lactam pharmacophore. Certain embodiments of these new inhibitors may also bind bacterial DD-peptidases, and thus may potentially act both as  $\beta$ -lactamase inhibitors and as antibiotic agents.

**[0014]** Li et al., Bioorg. Med. Chem. 5 (9):1783-1788 (1997), and Dryjanski and Pratt, Biochemistry 34:3569-3575 (1995), teach that  $\beta$ -lactamase enzymes are inactivated by 4-nitrophenyl *N*-(phenylmethylsulfonyl)aminomethylphosphonate and *p*-nitro-phenyl[(dansylamido)methyl]-posphonate, respectively. Despite the enzyme inhibitory activity of these compounds, however, neither compound is effective in producing a synergistic effect when co-administered with an antibiotic agent to a bacterially infected cell culture (see Table 1, compounds **1** and **72**). By contrast, the present inventors have surprisingly discovered that aryl- and (heteroaryl)-sulfonamidomethylphosphonate monoesters both inhibit enzymatic activity in vitro and enhance the potency of antibiotic agents in cell culture.

**[0015]** For purposes of the present invention, the following definitions will be used:

**[0016]** As used herein, the term " $\beta$ -lactamase inhibitor" is used to identify a compound having a structure as defined herein, which is capable of inhibiting  $\beta$ -lactamase activity. Inhibiting  $\beta$ -lactamase activity means inhibiting the activity of a class A, B, C, or D  $\beta$ -lactamase. Preferably, for antimicrobial applications such inhibition should be at a 50% inhibitory concentration below 100 micrograms/mL, more preferably below 30 micrograms/mL and most preferably below 10 micrograms/mL. The terms "class A", "class B", "class C", and "class D"  $\beta$ -lactamases are understood by those skilled in the art and can be found described in Waley, The Chemistry of  $\beta$ -Lactamase, Page Ed., Chapman & Hall, London, (1992) 198-228

**[0017]** In some embodiments of the invention, the  $\beta$ -lactamase inhibitor may also be capable of acting as an antibiotic agent by inhibiting bacterial cell-wall cross-linking enzymes. Thus, the term  $\beta$ -lactamase inhibitor is intended to encompass such dual-acting inhibitors. In certain preferred embodiments, the  $\beta$ -lactamase inhibitor may be capable of inhibiting D-alanyl-D-alanine-carboxypeptidases/transpeptidases (hereinafter DD-peptidases). The term "DD-peptidase" is used in its usual sense to denote penicillin-binding proteins involved in bacterial cell wall biosynthesis (see, e.g., Ghysen, Prospect. Biotechnol. 128:67-95 (1987)). In certain particularly preferred embodiments, the D-alanyl-D-alanine-carboxypeptidase/transpeptidase, which may be inhibited is the *Streptomyces* R61 DD-peptidase. This enzyme has conservation of active site mechanism with bacterial signal peptidases (see, e.g., Black et al., Current Pharmaceutical Design 4:133-154 (1998); Dalbey et al., Protein Science 6:1129-1138 (1997)). It is, therefore, possible that the  $\beta$ -lactamase inhibitors of the invention may also be capable of inhibition of bacterial signal peptidases

**[0018]** As used herein, the term "β-lactamase" denotes a protein capable of inactivating a β-lactam antibiotic. In one preferred embodiment, the β-lactamase is an enzyme which catalyzes the hydrolysis of the β-lactam ring of a β-lactam antibiotic. In certain preferred embodiments, the β-lactamase is microbial. In certain other preferred embodiments, the β-lactamase is a serine β-lactamase. Examples of such preferred β-lactamases are well known and are disclosed in, e.g., Waley, The Chemistry of β-Lactamase, Page Ed., Chapman & Hall, London, (1992) 198-228. In particularly preferred embodiments, the β-lactamase is a class C β-lactamase of *Enterobacter cloacae* P99 (hereinafter P99 β-lactamase), or a class A β-lactamase of the TEM-2 plasmid (hereinafter TEM β-lactamase).

**[0019]** As used herein, the term "organism" refers to any multicellular organism. Preferably, the organism is an animal, more preferably a mammal, and most preferably a human

**[0020]** For simplicity, chemical moieties are defined and referred to throughout primarily as univalent chemical moieties (e.g., alkyl, aryl, etc.). Nevertheless, such terms are also used to convey corresponding multivalent moieties under the appropriate structural circumstances clear to those skilled in the art. For example, while an "alkyl" moiety generally refers to a monovalent radical (e.g.  $\text{CH}_3\text{-CH}_2\text{-}$ ), in certain circumstances a bivalent linking moiety can be "alkyl," in which case those skilled in the art will understand the alkyl to be a divalent radical (e.g.,  $\text{-CH}_2\text{-CH}_2\text{-}$ ), which is equivalent to the term "alkylene." (Similarly, in circumstances in which a divalent moiety is required and is stated as being "aryl," those skilled in the art will understand that the term "aryl" refers to the corresponding divalent moiety, arylene.) All atoms are understood to have their normal number of valences for bond formation (i.e., 4 for carbon, 3 for N, 2 for O, and 2, 4, or 6 for S, depending on the oxidation state of the S). On occasion a moiety may be defined, for example, as  $(\text{A})_a\text{-B-}$ , wherein a is 0 or 1. In such instances, when a is 0 the moiety is B- and when a is 1 the moiety is A-B-. Also, a number of moieties disclosed herein exist in multiple tautomeric forms, all of which are intended to be encompassed by any given tautomeric structure.

**[0021]** The term "alkyl" as employed herein refers to straight and branched chain aliphatic groups having from 1 to 12 carbon atoms, preferably 1-8 carbon atoms, more preferably 1-6 carbon atoms, which is optionally substituted with one, two or three substituents. Unless otherwise specified, the alkyl group may be saturated, unsaturated, or partially unsaturated. As used herein, therefore, the term "alkyl" is specifically intended to include alkenyl and alkynyl groups, as well as saturated alkyl groups. Preferred alkyl groups include, without limitation, methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, isobutyl, pentyl, hexyl, vinyl, allyl, isobutenyl, ethynyl, and propynyl.

**[0022]** As employed herein, a "substituted" alkyl, cycloalkyl, aryl, or heterocyclic group is one having between one and about four, preferably between one and about three, more preferably one or

two, non-hydrogen substituents. Suitable substituents include, without limitation, halo, hydroxy, nitro, haloalkyl, alkyl, alkaryl, aryl, aralkyl, alkoxy, aryloxy, amino, acylamino, alkylcarbamoyl, arylcarbamoyl, aminoalkyl, alkoxy carbonyl, carboxy, hydroxyalkyl, alkanesulfonyl, arenesulfonyl, alkanesulfonamido, arenesulfonamido, aralkylsulfonamido, alkylcarbonyl, acyloxy, cyano, and ureido groups.

**[0023]** The term "cycloalkyl" as employed herein includes saturated and partially unsaturated cyclic hydrocarbon groups having 3 to 12, preferably 3 to 8 carbons, wherein the cycloalkyl group additionally is optionally substituted. Preferred cycloalkyl groups include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl.

**[0024]** An "aryl" group is a C<sub>6</sub>-C<sub>14</sub> aromatic moiety comprising one to three aromatic rings, which is optionally substituted. Preferably, the aryl group is a C<sub>6</sub>-C<sub>10</sub> aryl group. Preferred aryl groups include, without limitation, phenyl, naphthyl, anthracenyl, and fluorenyl. An "aralkyl" or "arylalkyl" group comprises an aryl group covalently linked to an alkyl group, either of which may independently be optionally substituted or unsubstituted. Preferably, the aralkyl group is C<sub>1-6</sub>alk(C<sub>6-10</sub>)aryl, including, without limitation, benzyl, phenethyl, and naphthylmethyl. An "alkaryl" or "alkylaryl" group is an aryl group having one or more alkyl substituents. Examples of alkaryl groups include, without limitation, tolyl, xylyl, mesityl, ethylphenyl, *tert*-butylphenyl, and methylnaphthyl.

**[0025]** A "heterocyclic" group is a ring structure having from about 3 to about 8 atoms, wherein one or more atoms are selected from the group consisting of N, O, and S. The heterocyclic group is optionally substituted on carbon with oxo or with one of the substituents listed above. The heterocyclic group may also independently be substituted on nitrogen with alkyl, aryl, aralkyl, alkylcarbonyl, alkylsulfonyl, arylcarbonyl, arylsulfonyl, alkoxy carbonyl, aralkoxy carbonyl, or on sulfur with oxo or lower alkyl. Preferred heterocyclic groups include, without limitation, epoxy, aziridinyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, piperazinyl, thiazolidinyl, oxazolidinyl, oxazolidinonyl, and morpholino.

**[0026]** In certain preferred embodiments, the heterocyclic group is a heteroaryl group. As used herein, the term "heteroaryl" refers to groups having 5 to 14 ring atoms, preferably 5, 6, 9, or 10 ring atoms; having 6, 10, or 14  $\pi$  electrons shared in a cyclic array; and having, in addition to carbon atoms, between one and about three heteroatoms selected from the group consisting of N, O, and S. Preferred heteroaryl groups include, without limitation, thienyl, benzothienyl, furyl, benzofuryl, dibenzofuryl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, indolyl, quinolyl, isoquinolyl, quinoxalinyl, tetrazolyl, oxazolyl, thiazolyl, and isoxazolyl.

**[0027]** In certain other preferred embodiments, the heterocyclic group is fused to an aryl or heteroaryl group. Examples of such fused heterocycles include, without limitation, tetrahydroquinolinyl and dihydrobenzofuranyl.

**[0028]** Additional preferred heterocyclis and heteroaryls include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolyl, decahydroquinolyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, 1H-indazolyl, indolenyl, indolyl, indolizyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolyl, isoindolyl, isothiazolyl, methylenedioxyphenyl, morpholyl, naphthyridinyl, octahydroisoquinolyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, quinazolyl, 4H-quinolizyl, quinuclidinyl, tetrahydroisoquinolyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

**[0029]** A moiety that is substituted is one in which one or more hydrogens have been independently replaced with another chemical substituent. As a non-limiting example, substituted phenyls include 2-fluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluoro-phenyl, 2-fluor-3-propylphenyl. As another non-limiting example, substituted n-octyls include 2,4 dimethyl-5-ethyl-octyl and 3-cyclopentyl-octyl. Included within this definition are methylenes (-CH<sub>2</sub>-) substituted with oxygen to form carbonyl (-CO-).

**[0030]** Unless otherwise stated, as employed herein, when a moiety (e.g., cycloalkyl, hydrocarbyl, aryl, heteroaryl, heterocyclic, úrea, etc.) is described as "optionally substituted" it is meant that the group optionally has from one to four, preferably from one to three, more preferably one or two, non-hydrogen substituents. Suitable substituents include, without limitation, halo, hydroxy, oxo (e.g., an annular -CH- substituted with oxo is -C(O)-) nitro, halohydrocarbyl, hydrocarbyl, aryl, aralkyl, alkoxy, aryloxy, amino, acylamino, alkylcarbamoyl, arylcarbamoyl, aminoalkyl, acyl, carboxy, hydroxyalkyl, , alkanesulfonyl, arenesulfonyl, alkanesulfonamido, arenesulfonamido, aralkylsulfonamido, alkylcarbonyl, acyloxy, cyano, and ureido groups. Preferred substituents, which are themselves not further substituted (unless expressly stated otherwise) are:

- (a) halo, cyano, oxo, carboxy, formyl, nitro, amino, amidino, guanidino,
- (b) C<sub>1</sub>-C<sub>5</sub> alkyl or alkenyl or arylalkyl imino, carbamoyl, azido, carboxamido, mercapto, hydroxy, hydroxyalkyl, alkylaryl, arylalkyl, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkenyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> alkoxycarbonyl, aryloxycarbonyl, C<sub>2</sub>-C<sub>8</sub> acyl, C<sub>2</sub>-C<sub>8</sub> acylamino, C<sub>1</sub>-C<sub>8</sub> alkylthio,

arylalkylthio, arylthio, C<sub>1</sub>-C<sub>8</sub> alkylsulfinyl, arylalkylsulfinyl, arylsulfinyl, C<sub>1</sub>-C<sub>8</sub> alkylsulfonyl, arylalkylsulfonyl, arylsulfonyl, C<sub>0</sub>-C<sub>6</sub> N-alkyl carbamoyl, C<sub>2</sub>-C<sub>15</sub> N,N-dialkylcarbamoyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, aroyl, aryloxy, arylalkyl ether, aryl, aryl fused to a cycloalkyl or heterocycle or another aryl ring, C<sub>3</sub>-C<sub>7</sub> heterocycle, or any of these rings fused or spiro-fused to a cycloalkyl, heterocyclyl, or aryl, wherein each of the foregoing is further optionally substituted with one more moieties listed in (a), above; and

- (c)  $-(CH_2)_s-NR^{30}R^{31}$ , wherein s is from 0 (in which case the nitrogen is directly bonded to the moiety that is substituted) to 6, and R<sup>30</sup> and R<sup>31</sup> are each independently hydrogen, cyano, oxo, carboxamido, amidino, C<sub>1</sub>-C<sub>8</sub> hydroxyalkyl, C<sub>1</sub>-C<sub>3</sub> alkylaryl, aryl-C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkenyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> alkoxycarbonyl, aryloxycarbonyl, aryl-C<sub>1</sub>-C<sub>3</sub> alkoxycarbonyl, C<sub>2</sub>-C<sub>8</sub> acyl, C<sub>1</sub>-C<sub>8</sub> alkylsulfonyl, arylalkylsulfonyl, arylsulfonyl, aroyl, aryl, cycloalkyl, heterocyclyl, or heteroaryl, wherein each of the foregoing is further optionally substituted with one more moieties listed in (a), above; or

R<sup>30</sup> and R<sup>31</sup> taken together with the N to which they are attached form a heterocyclyl or heteroaryl, each of which is optionally substituted with from 1 to 3 substituents from (a), above.

**[0031]** The term "halogen" or "halo" as employed herein refers to chlorine, bromine, fluorine, or iodine

**[0032]** As herein employed, the term "acyl" refers to an alkylcarbonyl or arylcarbonyl substituent.

**[0033]** The term "acylamino" refers to an amide group attached at the nitrogen atom. The term "carbamoyl" refers to an amide group attached at the carbonyl carbon atom. The nitrogen atom of an acylamino or carbamoyl substituent may be additionally substituted. The term "sulfonamido" refers to a sulfonamide substituent attached by either the sulfur or the nitrogen atom. The term "amino" is meant to include NH<sub>2</sub>, alkylamino, arylamino, and cyclic amino groups.

**[0034]** The term "ureido" as employed herein refers to a substituted or unsubstituted urea moiety.

**[0035]** The structure:




represents -S-, -S(O)- or -S(O)<sub>Z</sub>- when Z is S and n is 0, 1, or 2, respectively, -CH<sub>2</sub>- when Z is CH<sub>2</sub> and n is 0, and -C(O)- when Z is C and n is 1.

**[0036]** Preferred embodiments of a particular genus of compounds of the invention include combinations of preferred embodiments. For example, preferred R<sup>2</sup> substituents are given in paragraph [0046] and preferred R<sup>3</sup> substituents are given in paragraph [0047]. Thus, preferred compounds

according to the invention include those in which  $R^2$  is as described in paragraph [0046] and  $R^3$  is as described in paragraph [0047].

## Compounds

**[0037]** In a first aspect, the invention provides novel  $\beta$ -lactamase inhibitors. In one embodiment of the invention, the novel  $\beta$ -lactamase inhibitor is a compound of Formula (I):



or a pro-drug or pharmaceutically acceptable salt thereof, wherein

R<sup>1</sup> is aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted;

Z is C, CH<sub>2</sub>, or S;

n is 0, 1, or 2 when Z is S, n is 1 when Z is C, and n is 0 when Z is CH<sub>2</sub>;

L is C<sub>0</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>3</sub>alkyl-O-, -C(O)-, or -C(=N-OCH<sub>3</sub>);

R<sup>2</sup> is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, and aryl, any of which is optionally substituted;

R<sup>3</sup> is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, aryl, heteroaryl, and (heteroaryl)alkyl, any of which groups is optionally substituted;

$R^4$  is selected from the group consisting of OH, F,  $SR^7$ , and  $N(R^7)_2$ ; and

$R^5$  is selected from the group consisting of F,  $OR^6$ ,  $SR^7$ , and  $N(R^7)_2$ ,

R<sup>6</sup> is selected from the group consisting of -H, alkyl, cycloalkyl, C<sub>0</sub>-C<sub>3</sub>-alkyl-cycloalkyl, aryl, C<sub>0</sub>-C<sub>3</sub>-alkaryl, aralkyl, (heteroaryl)alkyl, heteroaryl, and C<sub>0</sub>-C<sub>3</sub>-alkyl-heteroaryl, wherein the aryl or heteroaryl portion of any such group is optionally substituted, and


R<sup>7</sup> at each occurrence is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, and aryl, each of which is optionally substituted;

provided that R<sup>4</sup> and R<sup>5</sup> are not both F, and further provided that R<sup>1</sup> is not 5-dimethylamino-1-naphthyl when R<sup>2</sup> and R<sup>3</sup> are both H, R<sup>4</sup> is OH, and R<sup>5</sup> is 4-nitrophenoxy.

**[0038]** In one preferred embodiment of the compound according to paragraph [0037], Z is S.

**[0039]** In another embodiment, the novel  $\beta$ -lactamase inhibitor is a compound of Formula (I):





I

or a pro-drug or pharmaceutically acceptable salt thereof, wherein

$R^1$  is aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted;

Z is C,  $CH_2$ , or S;

n is 0, 1, or 2 when Z is S, n is 1 when Z is C, and n is 0 when Z is  $CH_2$ ;

L is  $C_0$ - $C_3$ alkyl,  $C_1$ - $C_3$ alkyl-O-, -C(O)-, or -C(=N-OCH<sub>3</sub>)-;

$R^2$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, and aryl, wherein the aryl portion of any such group is optionally substituted;


$R^3$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, aryl, heteroaryl, and (heteroaryl)alkyl, wherein the aryl or heteroaryl portion of any such group is optionally substituted;

$R^4$  is  $OR^8$ , where  $R^8$  is selected from the group consisting of phenyl substituted with at least one chloro, nitro, or fluoro substituent; heteroaryl; and substituted heteroaryl; and

$R^5$  is  $OR^6$ , where  $R^6$  is selected from the group consisting of H, alkyl, cycloalkyl, aryl, aralkyl, (heteroaryl)alkyl, and heteroaryl, wherein the aryl or heteroaryl portion of any such group is optionally substituted.

**[0040]** In one preferred embodiment of the compound according to paragraph [0039], Z is S.

**[0041]** In yet another embodiment, the novel  $\beta$ -lactamase inhibitor is a compound of Formula (II):



II

or a pro-drug or pharmaceutically acceptable salt thereof, wherein

$R^1$  is aryl or heteroaryl, wherein the aryl or heteroaryl group is optionally substituted;

Z is C,  $CH_2$ , or S;

n is 0, 1, or 2 when Z is S, n is 1 when Z is C, and n is 0 when Z is  $CH_2$ ;

L is  $C_0$ - $C_3$ alkyl,  $C_1$ - $C_3$ alkyl-O-, -C(O)-, or -C(=N-OCH<sub>3</sub>)-;

Y is O,  $NR^7$ , or S;

$R^2$  is selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, and aryl;

$R^4$  and  $R^5$  are independently selected from the group consisting of OH, F,  $SR^7$ ,  $N(R^7)_2$ ,

and OR<sup>6</sup>,

where R<sup>6</sup> is selected from the group consisting of alkyl, cycloalkyl, aryl, aralkyl, (heteroaryl)alkyl, and heteroaryl, wherein the aryl or heteroaryl portion of any such group is optionally substituted, and where R<sup>7</sup> at each occurrence is independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, aralkyl, and aryl;

provided that R<sup>4</sup> and R<sup>5</sup> are not both F or both OH.

**[0042]** With respect to the compounds of Formula (I) or Formula (II), the following preferred values are applicable:

**[0043]** In certain preferred embodiments, R<sup>1</sup> is C<sub>6-14</sub>aryl, more preferably C<sub>6-10</sub>aryl, most preferably phenyl or naphthyl, any of which may be substituted. In certain other preferred embodiments, R<sup>1</sup> is heteroaryl or substituted heteroaryl. Preferably, the heteroaryl group is selected from the group consisting of thienyl, benzothienyl, furyl, benzofuryl, quinolyl, isoquinolyl, and thiazolyl. In certain particularly preferred embodiments, the heteroaryl group is thienyl or benzothienyl

**[0044]** Substituted aryl or heteroaryl groups have one or more, preferably between one and about three, more preferably one or two substituents, which are preferably selected from the group consisting of C<sub>1-6</sub>alkyl, preferably C<sub>1-4</sub>alkyl; halo, preferably Cl, Br, or F; haloalkyl, preferably (halo)<sub>1-5</sub>(C<sub>1-6</sub>)alkyl, more preferably (halo)<sub>1-5</sub>(C<sub>1-3</sub>)alkyl, and most preferably CF<sub>3</sub>; C<sub>1-6</sub>alkoxy, preferably methoxy, ethoxy, or benzyloxy; C<sub>6-10</sub>aryloxy, preferably phenoxy; C<sub>1-6</sub>alkoxycarbonyl, preferably C<sub>1-3</sub>alkoxycarbonyl, most preferably carbomethoxy or carboethoxy; C<sub>6-10</sub>aryl, preferably phenyl; (C<sub>6-10</sub>)ar(C<sub>1-6</sub>)alkyl, preferably (C<sub>6-10</sub>)ar(C<sub>1-3</sub>)alkyl, more preferably benzyl, naphthylmethyl or phenethyl; hydroxy(C<sub>1-6</sub>)alkyl, preferably hydroxy(C<sub>1-3</sub>)alkyl, more preferably hydroxymethyl; amino(C<sub>1-6</sub>)alkyl, preferably amino(C<sub>1-3</sub>)alkyl, more preferably aminomethyl; (C<sub>1-6</sub>)alkylamino, preferably methylamino, ethylamino, or propylamino; di(C<sub>1-6</sub>)alkylamino, preferably dimethylamino or diethylamino; (C<sub>1-6</sub>)alkylcarbamoyl, preferably methylcarbamoyl, dimethylcarbamoyl, or benzylcarbamoyl; (C<sub>6-10</sub>)arylcarbamoyl, preferably phenylcarbamoyl; (C<sub>1-6</sub>)alkaneacylamino, preferably acetylamino; (C<sub>6-10</sub>)areneacylamino, preferably benzoylamino; (C<sub>1-6</sub>)alkanesulfonyl, preferably methanesulfonyl; (C<sub>1-6</sub>)alkanesulfonamido, preferably methanesulfonamido; (C<sub>6-10</sub>)arenesulfonyl, preferably benzenesulfonyl or toluenesulfonyl; (C<sub>6-10</sub>)arenesulfonamido, preferably benzenesulfonyl or toluenesulfonyl; (C<sub>6-10</sub>)ar(C<sub>1-6</sub>)alkylsulfonamido, preferably benzylsulfonamido; C<sub>1-6</sub>alkylcarbonyl, preferably C<sub>1-3</sub>alkylcarbonyl, more preferably acetyl; (C<sub>1-6</sub>)acyloxy, preferably acetoxy; cyano; amino; carboxy; hydroxy; ureido; and nitro.

**[0045]** Preferably, n is 1 or 2. More preferably n is 2.

**[0046]**  $R^2$  is preferably selected from the group consisting of H;  $C_{1-8}$ alkyl, preferably  $C_{1-6}$ alkyl, more preferably  $C_{1-4}$ alkyl;  $C_{3-8}$ cycloalkyl, preferably cyclopropyl, cyclopentyl, or cyclohexyl;  $(C_{6-10})ar(C_{1-6})$ alkyl, preferably  $(C_{6-10})ar(C_{1-3})$ alkyl, more preferably benzyl; and  $C_{6-10}$ aryl, preferably phenyl; any of which groups is optionally substituted. More preferably,  $R^2$  is one of H, methyl, ethyl, propyl, cyclopropyl or benzyl. Most preferably,  $R^2$  is H.

**[0047]**  $R^3$  is preferably selected from the group consisting of H;  $C_{1-8}$ alkyl, preferably  $C_{1-6}$ alkyl, more preferably  $C_{1-4}$ alkyl;  $C_{3-8}$ cycloalkyl, preferably cyclopropyl, cyclopentyl, or cyclohexyl;  $(C_{6-10})ar(C_{1-6})$ alkyl, preferably  $(C_{6-10})ar(C_{1-3})$ alkyl, more preferably benzyl; heterocyclic having one or more, preferably between one and about three, more preferably one or two, ring atoms independently selected from the group consisting of N, O, and S; heterocyclic  $(C_{1-6})$ alkyl, preferably heterocyclic  $(C_{1-3})$ alkyl; and  $C_{6-10}$ aryl, preferably phenyl; any of which groups may be optionally substituted. More preferably,  $R^3$  is one of H, methyl, ethyl, propyl, isopropyl, isobutyl, butyl, phenyl, or benzyl.

**[0048]** In certain other preferred embodiments,  $R^2$  and  $R^3$  are taken together with the carbon and nitrogen atoms to which they are attached to form a nitrogen-containing heterocyclic ring. Preferably, the heterocyclic ring is a 4- to 7-membered ring, more preferably a 5- or 6-membered ring.

**[0049]** In certain preferred embodiments,  $R^4$  is preferably OH. In certain other preferred embodiments,  $R^4$  is preferably  $OR^8$ , wherein  $R^8$  is preferably selected from the group consisting of phenyl or heteroaryl, either of which may be optionally substituted. Where  $R^8$  is phenyl, the phenyl group is preferably substituted with at least one ester, amide, chloro, nitro, or fluoro substituent; more preferably with at least one chloro, nitro, or fluoro substituent; and most preferably with at least one nitro substituent.

**[0050]**  $R^5$  is preferably selected from the group consisting of F,  $OR^6$ ,  $SR^7$ , and  $N(R^7)_2$ , where  $R^6$  and  $R^7$  are as defined below. More preferably,  $R^5$  is F or  $OR^6$ . In certain particularly preferred embodiments,  $R^5$  is a good leaving group. Leaving group ability is understood by those skilled in the art, and is generally described in March, Advanced Organic Chemistry, Third Edition, John Wiley & Sons, 1985, pp 310-316. Compounds according to this embodiment of the invention are subject to attack by a nucleophilic residue such as serine on the enzyme, resulting in irreversible inactivation of the enzyme.

**[0051]**  $R^6$  is preferably selected from the group consisting of  $C_{1-8}$ alkyl, preferably  $C_{1-6}$ alkyl, more preferably  $C_{1-4}$ alkyl;  $C_{3-8}$ cycloalkyl, preferably cyclopropyl, cyclopentyl, or cyclohexyl;  $(C_{6-10})ar(C_{1-6})$ alkyl, preferably  $(C_{6-10})ar(C_{1-3})$ alkyl, more preferably benzyl; heterocyclic having one or more, preferably between one and about three, more preferably one or two, ring atoms independently selected from the group consisting of N, O, and S; heterocyclic  $(C_{1-6})$ alkyl, preferably heterocyclic  $(C_{1-3})$ alkyl; and  $C_{6-10}$ aryl,

preferably phenyl; any of which groups may be optionally substituted. Preferably, the heterocyclic group is heteroaryl.

**[0052]** More preferably,  $R^6$  is  $C_{6-10}$ aryl or heteroaryl, either of which is optionally substituted with between one and about three, more preferably one or two substituents, which are preferably selected from the group consisting of  $C_{1-6}$ alkyl, preferably  $C_{1-4}$ alkyl; halo, preferably Cl, Br, or F; haloalkyl, preferably (halo)<sub>1-5</sub>( $C_{1-6}$ )alkyl, more preferably (halo)<sub>1-5</sub>( $C_{1-3}$ )alkyl, and most preferably  $CF_3$ ;  $C_{1-6}$ alkoxy, preferably methoxy, ethoxy, or benzyloxy;  $C_{6-10}$ aryloxy, preferably phenoxy;  $C_{1-6}$ alkoxycarbonyl, preferably  $C_{1-3}$ alkoxycarbonyl, most preferably carbomethoxy or carboethoxy;  $C_{6-10}$ aryl, preferably phenyl; ( $C_{6-10}$ )ar( $C_{1-6}$ )alkyl, preferably ( $C_{6-10}$ )ar( $C_{1-3}$ )alkyl, more preferably benzyl, naphthylmethyl or phenethyl; hydroxy( $C_{1-6}$ )alkyl, preferably hydroxy( $C_{1-3}$ )alkyl, more preferably hydroxymethyl; amino( $C_{1-6}$ )alkyl, preferably amino( $C_{1-3}$ )alkyl, more preferably aminomethyl; ( $C_{1-6}$ )alkanesulfonyl, preferably methanesulfonyl; ( $C_{1-6}$ )alkanesulfonamido, preferably methanesulfonamido; ( $C_{6-10}$ )arenesulfonyl, preferably benzenesulfonyl or toluenesulfonyl; ( $C_{6-10}$ )arenesulfonamido, preferably benzenesulfonyl or toluenesulfonyl; ( $C_{6-10}$ )ar( $C_{1-6}$ )alkylsulfonamido, preferably benzylsulfonamido;  $C_{1-6}$ alkylcarbamoyl, preferably methylcarbamoyl, dimethylcarbamoyl, or benzylcarbamoyl;  $C_{6-10}$ arylcarbamoyl, preferably benzylcarbamoyl; ( $C_{1-6}$ )alkane-acylamino, preferably acetylamino; ( $C_{6-10}$ )areneacylamino, preferably benzoylamino;  $C_{1-6}$ alkylcarbonyl, preferably  $C_{1-3}$ alkylcarbonyl, more preferably acetyl; cyano; amino; carboxy; hydroxy; and nitro.

**[0053]** Most preferably,  $R^6$  is an aryl or heteroaryl group substituted with one or more substituents selected from the group consisting of halo, preferably chloro or fluoro; haloalkyl, preferably trifluoromethyl; nitro; cyano; acyl; carboxy; and alkoxycarbonyl. More preferably, the aryl or heteroaryl group has one or more chloro, fluoro, or nitro substituents. Most preferably,  $R^6$  is selected from the group consisting of nitrophenyl, pentafluorophenyl, trifluorophenyl, pyridyl, chloropyridyl, isoquinolyl, and quinolyl.

**[0054]**  $R^7$  is preferably selected from the group consisting of H;  $C_{1-8}$ alkyl, preferably  $C_{1-6}$ alkyl, more preferably  $C_{1-4}$ alkyl;  $C_{3-8}$ cycloalkyl, preferably cyclopropyl, cyclopentyl, or cyclohexyl; ( $C_{6-10}$ )ar( $C_{1-6}$ )alkyl, preferably ( $C_{6-10}$ )ar( $C_{1-3}$ )alkyl, more preferably benzyl; and  $C_{6-10}$ aryl, preferably phenyl; any of which groups may be optionally substituted.

**[0055]** With respect to the compounds of Formula (II), Y is O,  $NR^7$ , or S. Preferably, Y is  $NR^7$ , where  $R^7$  is as defined above. In certain particularly preferred embodiments, Y is NH.

**[0056]** In another preferred embodiment of the compound according to paragraph [0037], one or more of the following is true:

$R^1$  is aryl (preferably phenyl) or thienyl, each of which is optionally substituted;

L is methyl, methoxy, -C(O)-, or -C(=N-OCH<sub>3</sub>)-;

Z is C and n is 1;

R<sup>2</sup> is H;


R<sup>3</sup> is H;

R<sup>4</sup> is -OH;

R<sup>5</sup> is F, -OR<sup>6</sup>, or -SR<sup>6</sup>; and/or

R<sup>6</sup> is aryl or heteroaryl, each of which is optionally substituted.

**[0057]** In a preferred embodiment of the compound according to paragraph [0056], Z is C and n is 1; R<sup>2</sup> is H; R<sup>3</sup> is H; and R<sup>4</sup> is -OH:



**[0058]** In a preferred embodiment of the compound according to paragraph [0057],

R<sup>1</sup> is phenyl or thien-2-yl, each optionally substituted;

L is a covalent bond, -CH<sub>2</sub>O-, -C(O)-, or -C(=N-OCH<sub>3</sub>)-; and



















R<sup>5</sup> is -halo or -OR<sup>10</sup> wherein R<sup>10</sup> is phenyl, pyridinyl, or quinolinyl, each optionally substituted, provided that when L is -CH<sub>2</sub>O-, R<sup>5</sup> is not -F or *p*-nitrophenyl.














**[0059]** Preferred substituents on the compound according to paragraph [0058] are -NO<sub>2</sub>, -CO<sub>2</sub>H, and halo. Preferably R<sup>1</sup> is unsubstituted in the compound according to paragraph [0058].

**[0060]** In a preferred embodiment of the compound according to paragraph [0058], R<sup>5</sup> is selected from:

F				
			and	-H.


**[0061]** In a preferred embodiment, the compounds according to paragraph [0058] are those in which the combination of R<sup>1</sup>-L and R<sup>5</sup> are selected from the following (in which "PNP" is *p*-nitrophenyl):

R <sup>1</sup> -L-	R <sup>5</sup>
	PNP
	
	
	PNP
	
	
	
	PNP
	
	
	PNP

R <sup>1</sup> -L-	R <sup>5</sup>
	
	
	
	
	
	
and	
	-OH


**[0062]** In another embodiment, the compounds of the invention are those in which the phosphonate moiety of the compound according to paragraph [0057] is replaced with a thiophosphonate moiety, provided, however, that when R<sup>1</sup>-L- is benzyloxy, R<sup>5</sup> is not -O-PNP.

**[0063]** In another preferred embodiment of the compound according to paragraph [0037], R<sup>1</sup> is 4,7-dichlorobenzo[b]thiophen-2-yl; L is a covalent bond; Z is S; n is 2; R<sup>2</sup> and R<sup>3</sup> are -H; R<sup>4</sup> is -OH; and R<sup>5</sup> is -OR<sup>6</sup>:



wherein

$R^1$  is



$R^3$  is -H or  $-CO_2R^9$ , wherein  $R^9$  is  $-C_1-C_3$  alkyl;

$R^6$  is  $-L^1-A(L^2-B)_s$ , wherein

$L^1$  is  $C_0-C_3$  alkyl optionally mono- to per-halogenated;

A is  $C_3-C_6$  cycloalkyl, aryl, or heteroaryl;

$L^2$  is a covalent bond or  $(C_0-C_3\text{-hydrocarbyl})-X^1-(C_0-C_3\text{-hydrocarbyl})$ , wherein  $X^1$  is  $-C(O)-$ ,  $-NH-$ ,  $-NH-C(O)-$ ,  $-C(O)-NH-$ , or heteroaryl;


B is -H,  $C_3-C_6$  cycloalkyl, aryl, or heteroaryl; and

s is 0 or 1;


wherein when s is 0,  $(L^2-B)_s$  is -H or halo, and A and B are independently optionally substituted with 1-3 moieties independently selected from the group consisting of halo,  $-NO_2$ ,  $-CO_2H$ ,  $-CN$ ,  $-C(O)NH_2$ ,  $-SO_2NH_2$ , or  $-C_0-C_3\text{-hydrocarbyl}-Y-(C_1-C_3\text{-hydrocarbyl})$  wherein Y is a covalent bond,  $-O-C(O)-$ ,  $-C(O)-$ ,  $-O-$ ,  $-S-$ ,  $-SO_2-$ ,  $-C(O)NH-$ , or  $-NH-C(O)-$ ; and

each alkyl moiety is optionally mono- to per-halogenated.

**[0064]** In a preferred embodiment of the compound according to paragraph [0063],  $R^3$  is H and  $R^1$  is



**[0065]** In another preferred embodiment of the compound according to paragraph [0063],  $R^3$  is  $-CO_2Et$  and  $R^1$  is





**[0066]** In a preferred embodiment of the compound according to paragraph [0064],  $L^1$  is -O- and A is phenyl or pyridinyl (preferably pyridin-3-yl), each optionally substituted as stated in paragraph [0063].

**[0067]** In one preferred embodiment of the compound of paragraph [0066], s is 0.

**[0068]** In another preferred embodiment of the compound of paragraph [0066], s is 1 and  $L^2$  is -C(O)-, -C(O)NH-, -NH-, 1,2,4-oxadiazolyl, or 1,3,4-oxadiazolyl and B is phenyl, pyridinyl, cyclopropyl, or thienyl, wherein B is optionally substituted.


**[0069]** Preferred substituents on the A and B rings include -F, -Cl, -Br, -CO<sub>2</sub>H, -C(O)O-CH<sub>3</sub>, -CF<sub>3</sub>, -OCH<sub>3</sub>, -OCF<sub>3</sub>, -CH<sub>3</sub>, -CN, -C(O)NH<sub>2</sub>, -S-CF<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -NO<sub>2</sub>, -CF<sub>3</sub>CF<sub>3</sub>, -SO<sub>2</sub>CF<sub>3</sub>, -SO<sub>2</sub>CF<sub>3</sub>CF<sub>3</sub>, and -SO<sub>2</sub>NH<sub>2</sub>.


**[0070]** In a particularly preferred embodiment of the compound according to paragraph [0064] one or both of the following are true:





- A is selected from phenyl and pyridinyl;
- B is selected from phenyl, tetraazolyl, cyclopropyl, pyridinyl, and thienyl.

**[0071]** In one preferred embodiment according to paragraph [0065],  $R^6$  is phenyl or *p*-nitro phenyl.


**[0072]** In another particularly preferred embodiment of the compounds according to paragraph [0064], the compound is selected from those in which -O- $R^6$  is;








**[0073]** In another preferred embodiment of the compound according to paragraph [0037],  $R^1$  is optionally substituted benzo[b]thiophenyl; L is a covalent bond; Z is S; n is 2;  $R^2$  and  $R^3$  are -H;  $R^4$  is -OH; and  $R^5$  is -OR<sup>6</sup>:



wherein

$R^1$  is



optionally substituted with 1-3 moieties independently selected from the group consisting of -F, -Cl, -Br, -CO<sub>2</sub>H, -C(O)O-CH<sub>3</sub>, -CF<sub>3</sub>, -OCH<sub>3</sub>, -OCF<sub>3</sub>, -C<sub>1</sub>-C<sub>6</sub> alkyl, -CN, -C(O)NH<sub>2</sub>, -SCF<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -NO<sub>2</sub>, -CF<sub>3</sub>CF<sub>3</sub>, -SO<sub>2</sub>CF<sub>3</sub>, -SO<sub>2</sub>CF<sub>3</sub>CF<sub>3</sub>, and -SO<sub>2</sub>NH<sub>2</sub>;

$R^6$  is -L<sup>1</sup>A(L<sup>2</sup>B)<sub>s</sub>, wherein

L<sup>1</sup> is C<sub>0</sub>-C<sub>3</sub>alkyl optionally mono- to per-halogenated;

A is C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, aryl, or heteroaryl;

L<sup>2</sup> is a covalent bond or (C<sub>0</sub>-C<sub>3</sub>-hydrocarbyl)-X<sup>1</sup>-(C<sub>0</sub>-C<sub>3</sub>-hydrocarbyl), wherein X<sup>1</sup> is -C(O)-, -NH-, -NH-C(O)-, -C(O)-NH-, or heteroaryl;

B is -H, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, aryl, or heteroaryl; and


s is 0 or 1;

wherein when s is 0, (L<sup>2</sup>B)<sub>s</sub> is -H or halo, and A and B are independently optionally substituted with 1-3 moieties independently selected from the group consisting of halo, -CF<sub>3</sub>, -NO<sub>2</sub>, -CO<sub>2</sub>H, -CN, -C(O)NH<sub>2</sub>, -SO<sub>2</sub>NH<sub>2</sub>, or -C<sub>0</sub>-C<sub>3</sub>-hydrocarbyl-Y-(C<sub>1</sub>-C<sub>3</sub>-hydrocarbyl) wherein Y is a covalent bond, -O-C(O)-, -C(O)-, -O-, -S-, -SO<sub>2</sub>-, -C(O)-NH-, or -NH-C(O)-; and each alkyl moiety is optionally mono- to per-halogenated.

**[0074]** In a preferred embodiment of the compound according to paragraph [0073],  $R^6$  is phenyl optionally substituted with 1-3 moieties independently selected from the group consisting of halo,  $-CF_3$ ,  $-NO_2$ ,  $-CO_2H$ ,  $-CN$ ,  $-C(O)NH_2$ ,  $-SO_2NH_2$ , or  $-C_0-C_3$ hydrocarbyl-Y-( $C_1-C_3$ hydrocarbyl), wherein Y is a covalent bond,  $-O-C(O)-$ ,  $-C(O)-$ ,  $-O-$ ,  $-S-$ ,  $-SO_2-$ ,  $-C(O)NH-$ , or  $-NH-C(O)-$ , and each alkyl moiety is optionally mono- to per-halogenated.


**[0075]** In a preferred embodiment of the compound according to paragraph [0074],  $R^1$  is optionally substituted with 1 or 2 moieties independently selected from the group consisting of F, Cl, Br and  $C_1-C_6$  alkyl.

**[0076]** In a preferred embodiment of the compound according to paragraph [0075],  $R^1$  is




**[0077]** In another preferred embodiment of the compound according to paragraph [0074],  $R^6$  is phenyl optionally substituted with 1 or 2 moieties independently selected from the group consisting of halo,  $-CF_3$ , and CN.


**[0078]** In another preferred embodiment of the compound according to paragraph [0075], the compound is selected from those in which  $-OR^6$  is;




**[0079]** In another particularly preferred embodiment, the compound according to paragraph [0073] is




**[0080]** In another particularly preferred embodiment, the compound according to paragraph [0073] is




**[0081]** In another particularly preferred embodiment, the compound according to paragraph [0073] is




**[0082]** In another particularly preferred embodiment, the compound according to paragraph [0073] is




**[0083]** In another particularly preferred embodiment, the compound according to paragraph [0073] is



**[0084]** In another particularly preferred embodiment, the compound according to paragraph [0073] is



**[0085]** In another particularly preferred embodiment, the compound according to paragraph [0073] is



**[0086]** The present inventors have discovered a correlation between hydrophilicity of  $\beta$ -lactamase inhibitors (as measured by log P) and their biological efficacy (expressed as synergy as


described in the Examples). The correlation (see Figure 1) is described by a V-shaped curve, with its point intercepting the x-axis at a log P value of about -0.4 and its arms extending from -0.4 to +0.6. Based on this observed trend in structure-activity relationships, the  $\beta$ -lactamase inhibitors of the invention preferably have high negative or high positive log P values. Preferably, the log P for the inhibitor is  $\leq -0.6$  or  $\geq 0$ , more preferably  $\leq -1$  or  $\geq 0.2$ , still more preferably  $\leq -1.2$  or  $\geq 0.4$ , and most preferably  $\leq -1.4$  or  $\geq 0.6$ . Nikaido and Vaara, *Microbiological Reviews* **49**, 1-32 (1985), and Livermore, *Scad. J. Infect. Dis., Suppl.* **74**, 15-22 (1991), teach that hydrophilicity and cell permeability govern the behavior of antimicrobial agents

**[0087]** The compounds of Formula (I) are preferably monoacids ( $R^4 = OH$ ). In certain preferred embodiments, the  $\beta$ -lactamase inhibitor is a salt of the compound of Formula (I), the salt preferably being formed by treating the compound of Formula (I) with a base so as to remove the phosphonate hydrogen atom. Non-limiting examples of bases which may be used to deprotonate the compound of Formula (I) include sodium hydroxide, potassium hydroxide, ammonium hydroxide, sodium bicarbonate, potassium bicarbonate, sodium carbonate, and potassium carbonate. Preferably, the counterion thereby introduced is a pharmaceutically acceptable counterion, including without limitation sodium, magnesium, calcium, or ammonium.

**[0088]** Generally, the compounds of the invention can be routinely synthesized using techniques known to those skilled in the art in conjunction with the teachings herein. The compounds of Formula (I) wherein  $R^3 = H$  can be prepared in certain preferred embodiments according to the general synthetic route depicted in Scheme 1. Thus, Arbuzov reaction of bromomethylphthalimide with a phosphite such as triethylphosphite is preferably conducted at elevated temperature, e.g., 145 °C, in a solvent such as xylenes to afford the phthalimidomethylphosphonate **III**. Treatment of **III** with a hydrazine such as methylhydrazine in an alcoholic solvent such as methanol effects phthalimide cleavage to afford the aminomethylphosphonate **IV**. Treatment of **IV** with a sulfonyl chloride, sulfinyl chloride, or sulfenyl chloride of the general formula **V** in an organic solvent such as methylene chloride, and in the presence of a base such as triethylamine, provides the *N*-sulfonyl-, *N*-sulfinyl-, or *N*-sulfenyl-aminomethylphosphonate **VI**. Treatment of **VI** with a silyl halide such as trimethylsilyl bromide at room temperature in a solvent such as methylene chloride effects cleavage of the phosphonate ester to provide the phosphonic acid **VII**. In situ activation of **VII** with trichloroacetonitrile in pyridine, followed by treatment at 100° C with an aryl or heteroaryl alcohol, such as phenol or substituted phenol, affords an aryl or heteroaryl phosphonate. Treatment with an


aqueous base such as sodium bicarbonate then provides the sodium salt **VIII**, which corresponds to the compound of Formula (I), wherein  $R^2 = R^3 = H$ .

Scheme I




[0089] In certain other preferred embodiments, (sulfonamido)methylphosphonates of formula **XIV** may be prepared according to the procedures illustrated in Scheme 2. Thus, the sulfonyl chloride of formula **IX** is treated with ammonium hydroxide to produce the corresponding sulfonamide of formula **X**. Treatment of **X** with paraformaldehyde in the presence of a phosphite such as trimethyl phosphite affords the phosphonate diester of formula **XI**. Deprotection is effected by treatment of **XI** with a silyl halide such as trimethylsilyl bromide to produce **XII**, which may be converted to **XIV** by treatment with trichloroacetonitrile in pyridine, followed by treatment with an aryl or heteroaryl alcohol, as described above. Alternatively, treatment of **XII** with a chlorinating agent such as sulfuryl chloride or thionyl chloride, followed by treatment with an aryl or heteroaryl alcohol, affords the diester **XIII**, which is mono-deprotected by treatment with base to afford **VIII**.

Scheme 2



**[0090]** Compounds of Formula (I), wherein R<sup>3</sup> is not hydrogen, are synthesized according to the synthetic route depicted in Scheme 3. Thus, treatment of a sulfonamide **X** with an aldehyde in the presence of acetyl chloride and a phosphite such as diethylphosphite affords the α-substituted (sulfonamido)methylphosphonate ester **XV**. The remaining steps are performed analogously to those described for the methods according to Schemes 1 and 2 above to afford the salt **XVII**, which corresponds to the compound of Formula (I), wherein R<sup>3</sup> is not hydrogen.


Scheme 3



**[0091]** The compounds of Formula (I), wherein R<sup>4</sup> is F, are prepared according to the synthetic route outlined in Scheme 4. Thus, the phosphonic acid **VII** is treated with a chlorinating agent such as sulfuryl chloride or thionyl chloride to produce the dichlorophosphine oxide **XIX**, which, without isolation, is then treated with tetrabutylammonium fluoride. Treatment with base then affords the salt **XX**, which corresponds to the compound of Formula (I), wherein R<sup>4</sup> is F.




## Scheme 4



[0092] The compounds of Formula (I), wherein  $R^2$  is not hydrogen, may be prepared according to the synthetic route depicted in Scheme 5. Thus, the *N*-sulfonyl-, *N*-sulfinyl-, or *N*-sulfenyl-aminomethylphosphonate **VI** is treated with an alkyl halide in the presence of a base such as cesium carbonate to afford the *N*-alkylated derivative **XXI**. Deprotection and monoesterification as described above then provide the *N,N*-disubstituted compound **XXIII**.

## Scheme 5

**Pharmaceutical Compositions**

[0093] In a second aspect, the invention provides pharmaceutical compositions comprising a  $\beta$ -lactamase inhibitor of the invention (as described in paragraph [0037] - [0072]) and a pharmaceutically acceptable carrier or diluent.

[0094] As employed herein, the term "pro-drug" refers to pharmacologically acceptable derivatives, e.g., esters and amides, such that the resulting biotransformation product of the derivative is the active drug. Pro-drugs are known in the art and are described generally in, e.g., Goodman and Gilman's, "Biotransformation of Drugs", In The Pharmacological Basis of Therapeutics, 8th Ed., McGraw Hill, Int. Ed. 1992, p. 13-15, which is hereby incorporated by reference in its entirety. Compounds of the invention may be formulated by any method well known in the art and may be prepared for administration by any route, including, without limitation, parenteral, oral, sublingual, transdermal, topical, intranasal, intratracheal, or intrarectal. In certain particularly

preferred embodiments, compounds of the invention are administered intravenously in a hospital setting. In certain other embodiments, administration may be preferably by the oral route.

**[0095]** Accordingly, another aspect of the invention is a composition comprising a compound of the invention and a pharmaceutically acceptable carrier. The characteristics of the carrier will depend on the route of administration. As used herein, the term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue, or organism. Thus, compositions and methods according to the invention may contain, in addition to the inhibitor, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The pharmaceutical composition of the invention may also contain other active factors and/or agents which enhance the inhibition of  $\beta$ -lactamases and/or DD-peptidases.

### ***Inhibition of Bacterial Growth***

**[0096]** In a third aspect, the invention provides methods for inhibiting bacterial growth, such methods comprising administering to a bacterial cell culture, or to a bacterially infected cell culture, tissue, or organism, a  $\beta$ -lactamase inhibitor of Formula (I) or Formula (II) as defined for the first aspect of the invention (as described in paragraph [0037] - [0072]).

**[0097]** Preferably, the bacteria to be inhibited by administration of a  $\beta$ -lactamase inhibitor of the invention are bacteria that are resistant to  $\beta$ -lactam antibiotics. More preferably, the bacteria to be inhibited are  $\beta$ -lactamase positive strains that are highly resistant to  $\beta$ -lactam antibiotics. The terms "resistant" and "highly resistant" are well-understood by those of ordinary skill in the art (see, e.g., Payne *et al.*, *Antimicrobial Agents and Chemotherapy* 38:767-772 (1994); Hanaki *et al.*, *Antimicrobial Agents and Chemotherapy* 30:1120-1126 (1995)). Preferably, "highly resistant" bacterial strains are those against which the MIC of methicillin is  $> 100 \mu\text{g/mL}$ . Preferably, "slightly resistant" bacterial strains are those against which the MIC of methicillin is  $> 25 \mu\text{g/mL}$ .

**[0098]** The methods according to this aspect of the invention are useful for inhibiting bacterial growth in a variety of contexts. In certain preferred embodiments, the compound of the invention is administered to an experimental cell culture in vitro to prevent the growth of  $\beta$ -lactam resistant bacteria. In certain other preferred embodiments the compound of the invention is administered to an animal, including a human, to prevent the growth of  $\beta$ -lactam resistant bacteria in vivo. The

method according to this embodiment of the invention comprises administering a therapeutically effective amount of a  $\beta$ -lactamase inhibitor according to the invention for a therapeutically effective period of time to an animal, including a human. Preferably, the  $\beta$ -lactamase inhibitor is administered in the form of a pharmaceutical composition according to the second aspect of the invention.

**[0099]** The terms “therapeutically effective amount” and “therapeutically effective period of time” are used to denote known treatments at dosages and for periods of time effective to show a meaningful patient benefit, *i.e.*, healing of conditions associated with bacterial infection, and/or bacterial drug resistance. Preferably, such administration should be parenteral, oral, sublingual, transdermal, topical, intranasal, intratracheal, or intrarectal. When administered systemically, the therapeutic composition is preferably administered at a sufficient dosage to attain a blood level of inhibitor of at least about 100 micrograms/mL, more preferably about 1 milligrams/mL, and still more preferably about 10 milligrams/mL. For localized administration, much lower concentrations than this may be effective, and much higher concentrations may be tolerated.

**[00100]** In certain preferred embodiments of the method according to this aspect of the invention, a  $\beta$ -lactamase inhibitor according to the invention is co-administered with an antibiotic. Preferably, such co-administration produces a synergistic effect. As employed herein, the terms “synergy” and “synergistic effect” indicate that the effect produced when two or more drugs are co-administered is greater than would be predicted based on the effect produced when the compounds are administered individually. While not wishing to be bound by theory, the present inventors believe that the  $\beta$ -lactamase inhibitors according to the invention act to prevent degradation of  $\beta$ -lactam antibiotics, thereby enhancing their efficacy and producing a synergistic effect. In particularly preferred embodiments of the invention, therefore, the co-administered antibiotic is a  $\beta$ -lactam antibiotic. For purposes of this invention, the term “co-administered” is used to denote simultaneous or sequential administration.

**[00101]** Synergy may be expressed as a ratio of the minimum inhibitory concentration (MIC) of an antibiotic tested in the absence of a  $\beta$ -lactamase inhibitor to the MIC of the same antibiotic tested in the presence of the  $\beta$ -lactamase inhibitor. A ratio of one (1) indicates that the  $\beta$ -lactamase inhibitor has no effect on antibiotic potency. A ratio greater than one (1) indicates that the  $\beta$ -lactamase inhibitor produces a synergistic effect when co-administered with the antibiotic agent. Preferably the  $\beta$ -lactamase inhibitor produces a synergy ratio of at least about 2, more preferably about 4, and still


more preferably about 8. Most preferably, the  $\beta$ -lactamase inhibitor produces a synergy ratio of at least about 16.

**[00102]** In certain other preferred embodiments, the  $\beta$ -lactamase inhibitor according to the invention may itself have antibiotic activity, and thus potentially can be administered alone or can be co-administered with a  $\beta$ -lactam antibiotic or any other type of antibiotic.

**[00103]** The term "antibiotic" is used herein to describe a compound or composition which decreases the viability of a microorganism, or which inhibits the growth or reproduction of a microorganism. "Inhibits the growth or reproduction" means increasing the generation cycle time by at least 2-fold, preferably at least 10-fold, more preferably at least 100-fold, and most preferably indefinitely, as in total cell death. As used in this disclosure, an antibiotic is further intended to include an antimicrobial, bacteriostatic, or bactericidal agent. Non-limiting examples of antibiotics useful according to this aspect of the invention include penicillins, cephalosporins, aminoglycosides, sulfonamides, macrolides, tetracyclins, lincosides, quinolones, chloramphenicol, vancomycin, metronidazole, rifampin, isoniazid, spectinomycin, trimethoprim, sulfamethoxazole, and others. The term " $\beta$ -lactam antibiotic" is used to designate compounds with antibiotic properties containing a  $\beta$ -lactam functionality. Non-limiting examples of  $\beta$ -lactam antibiotics useful according to this aspect of the invention include penicillins, cephalosporins, penems, carbapenems, and monobactams.

**[00104]** The following examples are intended to further illustrate certain preferred embodiments of the invention, and are not intended to limit the scope of the invention.

**Example 1: [(4-fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt. (30)**



Step 1. Dimethyl (phthalimidomethyl)phosphonate (ii).

**[00105]** A mixture of *N*-(bromomethyl) phthalimide (i) (10.43 g, 43.46 mmol) and trimethyl phosphite (5.93 g, 47.80 mmol) was heated at reflux in xylene (20 mL) for 6 h. The reaction mixture was then cooled to room temperature and concentrated. Crystallization from CHCl<sub>3</sub>/hexane gave (ii) (7.60 g, 65%) as a white solid: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 3.84 (d, *J*<sub>H,P</sub> = 10.8 Hz, 6 H), 4.12 (d, *J* = 11.4 Hz, 2 H), 7.76 (m, 2 H), 7.87 (m, 2 H).

Step 2. [(4-fluorobenzenesulfonylamino)methyl]-phosphonic acid dimethyl ester (35).

**[00106]** A solution of phthalimide (ii) (1.50 g, 5.66 mmol) in anhydrous CH<sub>3</sub>OH (15 mL) was treated with hydrazide monohydrate (0.29 mL, 6.0 mmol) and stirred at room temperature for 3.5 days. The white phthalyl hydrazide precipitate was filtered off and the filtrate was concentrated at temperature below 20 °C to give dimethyl aminomethyl phosphonate (iii) as a pale yellow oil. Without purification, compound (iii) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL), cooled to 0 °C, and treated with *N*-methylmorpholine (0.84 mL, 7.36 mmol) and *p*-fluorobenzenesulfonyl chloride (1.43 g, 7.36 mmol). The mixture was warmed to room temperature over 2 hours, and stirred for 16 hours. The reaction mixture was then diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) washed with 1 N HCl (2 × 25 mL), dried (MgSO<sub>4</sub>), filtered, and concentrated. Purification by flash chromatography (silica gel; elution with 4% CH<sub>3</sub>OH in EtOAc) gave **35** (0.58 g, 35%) as a white solid: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 3.29 (d, *J* = 13.2 Hz, 2 H), 3.78 (d, *J*<sub>H,P</sub> = 11.1 Hz, 6 H), 6.50 (br s, 1 H), 7.20 (m, 2 H), 7.90 (m, 2 H).

Step 3. [(4-fluorobenzenesulfonylamino)methyl]-phosphonic acid (iv).

**[00107]** A solution of diester **35** (0.58 g, 1.95 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and bromotrimethylsilane (1.5 mL, 11.7 mmol) was stirred at room temperature for 5 hours and then concentrated. The resulting residue was dissolved in anhydrous CH<sub>3</sub>OH (10 mL) and stirred at room temperature for 20 minutes. Insolubles were filtered, and the filtrate was concentrated. Purification by trituration with CH<sub>2</sub>Cl<sub>2</sub> gave the phosphonic acid (iv) (0.51 g, 97%) as a white solid: <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.17 (d, *J* = 13.5 Hz, 2 H), 7.35 (m, 2 H), 7.97 (m, 2 H).

Step 4. [(4-fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (30).

**[00108]** A mixture of compound (iv) (52.6 mg, 0.20 mmol), 2-chloro-3-hydroxypyridine (28.5 mg, 0.22 mmol), and CCl<sub>3</sub>CN (100 μL, 1.00 mmol) was heated at 105 °C in pyridine. After 6 h, the reaction mixture was cooled to room temperature and then concentrated. The residue was dissolved

in H<sub>2</sub>O (10 mL) containing 1 N NaHCO<sub>3</sub> (0.6 mL), washed with EtOAc (5 mL × 2) and the water layer was lyophilized. Purification by reverse-phase preparative TLC (C<sub>18</sub>-silica gel, 20% CH<sub>3</sub>CN in H<sub>2</sub>O) gave sodium monophosphonate salt **30** (30 mg, 38%) as a white fluffy solid: <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ<sub>H</sub> 3.17 (d, *J* = 12.6 Hz, 2 H), 7.16 (m, 2 H), 7.24 (dd, *J* = 4.8, 8.1 Hz, 1 H), 7.57 (m, 1 H), 7.77 (m, 2 H), 7.98 (m, 1 H); <sup>31</sup>P NMR (121 MHz, CD<sub>3</sub>OD) δ<sub>P</sub> 13.4.

**Example 2: [(Benzylsulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (1).**

[00109] Following procedures analogous to those described in Example 1, substituting benzylsulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained: <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O), δ<sub>H</sub> 3.23 (d, *J* = 11.7 Hz, 2 H), 4.40 (s, 2 H), 7.18 (d, *J* = 9.3 Hz, 2 H), 7.31 (m, 5 H), 8.13 (d, *J* = 9.3 Hz, 2 H); <sup>31</sup>P NMR (121 MHz, D<sub>2</sub>O) δ<sub>P</sub> 14.8.

**Example 3: [(4-Fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (2).**

[00110] Following procedures analogous to those described in Example 1, substituting 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained: <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.15 (d, *J* = 13.5 Hz, 2 H), 7.31 (m, 2 H), 7.40 (d, *J* = 9.3 Hz, 2 H), 7.95 (m, 2 H), 8.21 (d, *J* = 9.3 Hz, 2 H); <sup>31</sup>P NMR (121 MHz, CD<sub>3</sub>OD) δ<sub>P</sub> 12.4; <sup>13</sup>C NMR (75.4 MHz, CD<sub>3</sub>OD) δ<sub>C</sub> 40.8 (d, *J*<sub>C,P</sub> = 153 Hz), 117.2 (d, *J* = 22.69 Hz), 122.4 (d, *J* = 4.6 Hz), 126.2, 131.2 (d, *J* = 9.4 Hz), 137.1 (d, *J* = 3.2 Hz), 144.8, 159.4 (d, *J* = 7.7 Hz), 166.5 (d, *J*<sub>C,F</sub> = 252 Hz).

**Example 4: [(4-Chlorobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (3).**

[00111] Following procedures analogous to those described in Example 1, substituting 4-chlorobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained: <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ<sub>H</sub> 8.06 (d, *J* = 9.3 Hz, 2 H), 7.66 (d, *J* = 9.0 Hz, 2 H), 7.40 (d, *J* = 9.0 Hz, 2 H), 7.04 (d, *J* = 9.3 Hz, 2 H), 3.14 (d, *J* = 12.6 Hz, 2 H); <sup>31</sup>P NMR (121 MHz, D<sub>2</sub>O) δ<sub>P</sub> 14.1.

**Example 5: *[(2-naphthylsulfonylamino)methyl]-phosphonic acid mono-(4-nitro phenyl) ester sodium salt (4).***

**[00112]** Following procedures analogous to those described in Example 1, substituting 2-naphthylsulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.19 (br.s, 1 H), 7.61 (d,  $J = 9.3$  Hz, 2 H), 7.41-7.82 (m, 6 H), 6.73 (d,  $J = 9.3$  Hz, 2 H), 3.22 (d,  $J = 12.6$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  13.4.

**Example 6: *[(4-Methoxybenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (5).***

**[00113]** Following procedures analogous to those described in Example 1, substituting 4-methoxybenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.0 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.6 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.0 (d,  $J = 9$  Hz, 2 H, Ar-H), 6.8 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.8 (s, 3 H,  $\text{CH}_3$ ), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ ).

**Example 7: *[(4-Toluylsulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (6).***

**[00114]** Following procedures analogous to those described in Example 1, substituting 4-toluylsulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.1 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.6 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.0 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ ); 2.2 (s, 3 H,  $\text{CH}_3$ ).

**Example 8: *(Benzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitro-phenyl) ester sodium salt (7).***

**[00115]** Following procedures analogous to those described in Example 1, substituting benzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.1 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.4-7.8 (m, 5 H, Ar-H), 7.0 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ ).

**Example 9: *[(4-Fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(3-pyridyl) ester sodium salt (8).***

[00116] Following procedures analogous to those described in Example 1, substituting 3-hydroxypyridine for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.10-8.18 (m, 8 H), 3.08 (d,  $J = 12.6$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.7.

**Example 10: *[(3,4-Dichlorobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (9).***

[00117] Following procedures analogous to those described in Example 1, substituting 3,4-dichlorobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.24 (d,  $J = 9.3$  Hz, 2 H), 8.01 (d,  $J = 2.1$  Hz, 1 H), 7.74 (d,  $J = 2.1$  Hz, 1 H), 7.69 (s, 1 H), 7.21 (d,  $J = 9.3$  Hz, 2 H), 3.39 (d,  $J = 12.3$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.2.

**Example 11: *[(1-Naphthylsulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (10).***

[00118] Following procedures analogous to those described in Example 1, substituting 1-naphthylsulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.4 (d, 1 H, Ar-H), 8.2 (d, 1 H, Ar-H), 8.1 (d, 1 H, Ar-H), 7.8 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.4-7.6 (m, 3 H, Ar-H), 6.6 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  16.5 (t, 1 P).

**Example 12: *[(4-Fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(8-quinoliny) ester sodium salt (11).***

[00119] Following procedures analogous to those described in Example 1, substituting 8-hydroxyquinoline for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.69 (m, 1 H), 8.22 (m, 1 H), 6.90-7.89 (m, 8 H), 3.10 (d,  $J = 13.8$  Hz, 2 H).

**Example 13: *[(4-Nitrobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (12).***

[00120] Following procedures analogous to those described in Example 1, substituting 4-nitrobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-



chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.6 (d, 2 H, Ar-H), 8.4 (d,  $J=9$  Hz, 2 H, Ar-H), 8.0 (d, 2 H, Ar-H), 7.4 (d,  $J=9$  Hz, 2 H, Ar-H), 3.2 (d,  $J=13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  17.5 (t, 1 P).

**Example 14: *[(2-Nitrobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (13).***

[00121] Following procedures analogous to those described in Example 1, substituting 2-nitrobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.0 (d,  $J=9$  Hz, 2 H, Ar-H), 7.8 (s, 1 H, Ar-H), 7.6 (s, 1 H, Ar-H), 7.6 (d, 2 H, Ar-H), 7.1 (d,  $J=9$  Hz, 2 H, Ar-H), 3.2 (d,  $J=13$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  17.5 (t, 1 P).

**Example 15: *[(2,5-Dichlorobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (14).***

[00122] Following procedures analogous to those described in Example 1, substituting 2,5-dichlorobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.1 (d,  $J=9$  Hz, 2 H, Ar-H), 7.8 (s, 1 H, Ar-H), 7.4 (s, 2 H, Ar-H), 7.1 (d,  $J=9$  Hz, Ar-H), 3.2 (d,  $J=13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  17.5 (t, 1 P).

**Example 16: *[(Thiophene-2-sulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (15).***

[00123] Following procedures analogous to those described in Example 1, substituting 2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.09 (d,  $J=9.6$  Hz, 2 H), 7.69 (dd,  $J=5.1; 1.5$  Hz, 1 H), 7.57 (dd,  $J=3.6; 1.2$  Hz, 1 H), 7.12 (d,  $J=9.6$  Hz, 2 H), 7.05 (dd,  $J=5.1; 3.6$  Hz, 1 H), 3.16 (d,  $J=12.9$  Hz, 2 H).

**Example 17: *[(4-tert-Butylbenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (16).***

[00124] Following procedures analogous to those described in Example 1, substituting 4-tert-butylbenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-

chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.8 (m, 2 H, Ar-H), 7.6 (m, 2 H, Ar-H), 7.1 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ ), 1.2 (s, 9 H, t-Bu);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  18 (t, 1P).

**Example 18: [(4-Trifluoromethylbenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (17).**

[00125] Following procedures analogous to those described in Example 1, substituting 4-trifluoromethylbenzenesulfonyl for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.8 (m, 2 H, Ar-H), 7.6 (m, 2 H, Ar-H), 7.1 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  18 (t, 1P).

**Example 19: [(2,4-Dinitrobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (18).**

[00126] Following procedures analogous to those described in Example 1, substituting 2,4-dinitrobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.61 (m, 1 H); 8.35 (m, 1 H), 8.15 (m, 1 H), 7.98 (d,  $J = 9.0$  Hz, 2 H); 7.02 (d,  $J = 9.0$  Hz, 2 H), 3.41 (d,  $J = 11.4$  Hz, 2 H).

**Example 20: [(8-Quinolylsulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (19).**

[00127] Following procedures analogous to those described in Example 1, substituting 8-quinolylsulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  6.68-8.78 (m, 10 H), 3.13 (d,  $J = 12.9$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  13.4.

**Example 21: [2,4,6-trimethylbenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenol) ester sodium salt (21).**

[00128] Following procedures analogous to those described in Example 1, substituting 2,4,6-trimethylbenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$

8.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 6.8 (s, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ ), 2.3 (s, 6 H,  $\text{CH}_3$ ), 2.1 (s, 3 H,  $\text{CH}_3$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  18 (t, 1P).

**Example 22: [(4-Chloro-3-nitrobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (22).**

[00129] Following procedures analogous to those described in Example 1, substituting 4-chloro-3-nitrobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.4 (s, 1 H, Ar-H), 8.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.9 (d, 1 H, Ar-H), 7.7 (d, 1 H, Ar-H), 7.2 (s, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  18 (t, 1 P).

**Example 23: [2-Bromobenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (23).**

[00130] Following procedures analogous to those described in Example 1, substituting 2-bromobenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta$  8.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 8.1 (s, 1 H, Ar-H), 7.6 (d, 1 H, Ar-H), 7.4 (m, 2 H, Ar-H), 7.2 (s, 2 H, Ar-H), 3.2 (d,  $J = 13$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  17 (t, 1P).

**Example 24: [(3-Pyridinesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (24)**

[00131] Following procedures analogous to those described in Example 1, substituting 3-pyridinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.80 (s, 1 H), 8.55 (d,  $J = 4.8$  Hz, 1 H), 8.13 (d,  $J = 8.1$  Hz, 1 H), 8.03 (d,  $J = 9.0$  Hz, 2 H), 7.46 (dd,  $J = 4.8$ , 8.1 Hz, 1 H), 7.05 (d,  $J = 9.0$  Hz, 2 H), 3.19 (d,  $J = 12.3$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  13.7.

**Example 25: [(3-Pyridinesulfonylamino)methyl]-phosphonic acid mono-(3-pyridinyl) ester sodium salt (25).**

[00132] Following procedures analogous to those described in Example 1, substituting 3-pyridinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 3-hydroxypyridine for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$

3.17 (d,  $J = 13.2$  Hz, 2 H), 7.37 (dd,  $J = 4.8, 8.1$  Hz, 1 H), 7.57-7.69 (m, 2 H), 8.23-8.29 (m, 2 H), 8.40 (br s, 1 H), 8.74 (dd,  $J = 1.2, 4.8$  Hz, 1 H), 9.00 (d,  $J = 1.5$  Hz, 1 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  12.1.;  $^{13}\text{C}$  NMR (75.4 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  40.6 (d,  $J = 152$  Hz), 125.5, 125.7, 130.5 (d,  $J = 3.8$  Hz), 136.8, 138.2, 143.5 (d,  $J = 4.3$  Hz), 144.7, 148.7, 151.2 (d,  $J = 7.7$  Hz), 153.8.

**Example 26: [(3-Dibenzofuransulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (27).**

**[00133]** Following procedures analogous to those described in Example 1, substituting 3-dibenzofuransulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.2 (s, 1 H, Ar-H), 7.8 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.45 (m, 3 H, Ar-H), 7.2 (t,  $J = 7.5$  Hz, 2 H, Ar-H), 6.8 (d,  $J = 9$  Hz, 2 H, Ar-H), 3.2 (d,  $J = 12.9$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  17.17 (t,  $J = 12.2$  Hz, 1 P).

**Example 27: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(2,4,6-trifluorophenyl) ester sodium salt (28).**

**[00134]** Following procedures analogous to those described in Example 1, substituting 2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 2,4,6-trifluorophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ ),  $\delta_{\text{H}}$  7.73 (dd,  $J = 5.0; 1.5$  Hz, 1 H), 7.62 (dd,  $J = 3.6, 1.5$  Hz, 1 H), 7.10 (dd,  $J = 5.0, 3.6$  Hz, 1 H), 6.77 (m, 2 H), 3.20 (d,  $J = 13.2$  Hz, 2 H).

**Example 28: (2-Pyridinesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (31).**

**[00135]** Following procedures analogous to those described in Example 1, substituting 2-pyridinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ ),  $\delta_{\text{H}}$  8.49 (m, 1 H), 8.08 (d,  $J = 9.0$  Hz, 2 H), 7.86-7.98 (m, 2 H), 7.52 (m, 1 H), 7.11 (d,  $J = 9.0$  Hz, 2 H), 3.28 (d,  $J = 12.3$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ ),  $\delta_{\text{P}}$  13.7.

**Example 29: [(2-Pyridinesulfonylamino)methyl]-phosphonic acid mono-(2-chloro-pyridine-3-yl) ester sodium salt (32)**

[00136] Following procedures analogous to those described in Example 1, substituting 2-pyridinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.48 (d,  $J = 4.8$  Hz, 1 H), 7.83-7.98 (m, 3 H), 7.57 (d,  $J = 8.1$  Hz, 1 H), 7.51 (m, 1 H), 7.23 (dd,  $J = 4.8, 8.1$  Hz, 1 H), 3.30 (d,  $J = 12.0$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  19.0.

**Example 30: [(5-Isoquinolinesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (33).**

[00137] Following procedures analogous to those described in Example 1, substituting 5-isoquinolinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.92 (br s, 1 H), 8.31 (br s, 1 H), 8.21 (dd,  $J = 1.0, 7.5$  Hz, 1 H), 8.12 (d,  $J = 6.3$  Hz, 1 H), 7.98 (d,  $J = 8.1$  Hz, 1 H), 7.62 (d,  $J = 9.0$  Hz, 2 H), 7.52 (m, 1 H), 6.53 (d,  $J = 9.0$  Hz, 2 H), 3.18 (d,  $J = 12.9$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  12.5.

**Example 31: [(2-Pyridinesulfonylamino)methyl]-phosphonic acid (34).**

[00138] Following procedures analogous to those described in Example 1, steps 2-3, substituting 2-pyridinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.53 (m, 1 H), 7.90-8.05 (m, 2 H), 7.58 (m, 1 H), 3.07 (d,  $J = 12.6$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  15.8.

**Example 32: [(5-Isoquinolinesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (36).**

[00139] Following procedures analogous to those described in Example 1, substituting 5-isoquinolinesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  9.00 (s, 1 H), 8.36 (d,  $J = 6.0$  Hz, 1 H), 8.29 (d,  $J = 7.2$  Hz, 1 H), 8.19 (d,  $J = 6.0$  Hz, 1 H), 8.06 (d,  $J = 7.5$  Hz, 1 H), 7.74 (d,  $J = 4.8$  Hz, 1 H), 7.57 (m, 1 H), 7.07 (d,  $J = 8.1$  Hz, 1 H), 6.92 (dd,  $J = 4.8, 8.1$  Hz, 1 H), 3.24 (d,  $J = 11.7$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  12.1.

**Example 33: [(4-Fluorobenzenesulfonylamino)methyl]-phosphonic acid mono-(2-bromopyridin-3-yl) ester sodium salt (37).**

[00140] Following procedures analogous to those described in Example 1, substituting 2-bromo-3-hydroxypyridine for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.95 (d,  $J = 4.8$  Hz, 2 H), 7.76 (m, 2 H), 7.51 (d,  $J = 8.1$  Hz, 2 H), 7.25 (dd,  $J = 4.8, 8.1$  Hz, 1 H), 7.16 (m, 2 H), 3.17 (d,  $J = 12.9$  Hz, 2 H).

**Example 34: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (38).**

[00141] Following procedures analogous to those described in Example 1, substituting 2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.99 (d,  $J = 4.8$  Hz, 1 H), 7.70 (d,  $J = 4.8$  Hz, 1 H), 7.58 (m, 2 H), 7.26 (dd,  $J = 8.4; 5.1$  Hz, 1 H), 7.06 (t,  $J = 4.8$  Hz, 1 H), 3.21 (d,  $J = 12.9$  Hz, 2 H).

**Example 35: [(4-phenylbenzenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (39).**

[00142] Following procedures analogous to those described in Example 1, substituting 4-phenylbenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-nitrophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.87 (d,  $J = 9.0$  Hz, 2 H), 7.73 (d,  $J = 8.1$  Hz, 2 H), 7.59 (d,  $J = 8.1$  Hz, 2 H), 7.49 (d,  $J = 6.6$  Hz, 2 H), 7.35 (m, 3 H), 6.91 (d,  $J = 9.0$  Hz, 2 H), 3.16 (d,  $J = 12.3$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  12.6.

**Example 36: [(4-Phenylbenzenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridine-3-yl) ester sodium salt (40).**

[00143] Following procedures analogous to those described in Example 1, substituting 4-phenylbenzenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.04 (m, 1 H), 7.78 (d,  $J = 8.4$  Hz, 2 H), 7.65 (d,  $J = 8.4$  Hz, 2 H), 7.54 (d,  $J = 6.6$  Hz, 2 H), 7.31-7.43 (m, 3 H), 7.16 (m, 1 H), 3.14 (d,  $J = 12.6$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.3.

**Example 37: [(5-Bromo-2-thiophenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (41).**

[00144] Following procedures analogous to those described in Example 1, substituting 5-bromo-2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.10 (d,  $J = 8.4$  Hz, 2 H), 7.30 (d,  $J = 4.2$  Hz, 1 H), 7.10 (d,  $J = 8.4$  Hz, 2 H), 7.02 (d,  $J = 4.2$  Hz, 1 H), 3.18 (d,  $J = 12.9$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  13.3.

**Example 38: [(5-Bromo-2-thiophenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (42).**

[00145] Following procedures analogous to those described in Example 1, substituting 5-bromo-2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.96 (d,  $J = 4.8$  Hz, 1 H), 7.55 (d,  $J = 8.1$  Hz, 1 H), 7.31 (d,  $J = 3.9$  Hz, 1 H), 7.22 (dd,  $J = 4.8, 8.1$  Hz, 1 H), 7.01 (d,  $J = 3.9$  Hz, 1 H), 3.21 (d,  $J = 12.9$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  15.1.

**Example 39: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid (43).**

[00146] Following procedures analogous to those described in Example 1, steps 2-3, substituting 2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.79 (dd,  $J = 4.8, 12.6$  Hz, 1 H), 7.64 (dd,  $J = 3.9, 1.2$  Hz, 1 H), 7.17 (dd,  $J = 4.8, 3.9$  Hz, 1 H), 3.16 (d,  $J = 13.2$  Hz, 2 H).

**Example 40: [(5-Bromo-2-thiophenesulfonylamino)methyl]-phosphonic acid (58).**

[00147] Following procedures analogous to those described in Example 1, steps 2-3, substituting 5-bromo-2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.47 (d,  $J = 3.9$  Hz, 1 H), 7.25 (d,  $J = 3.9$  Hz, 1 H), 3.23 (d,  $J = 13.5$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  17.9;  $^{13}\text{C}$  NMR (75.4 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  40.8 (d,  $J = 157$  Hz), 120.3, 132.1, 133.7, 143.0.

**Example 41: [(3-(N-Phenylcarbamoylamino)-sulfonylamino)methyl]-phosphonic acid (59).**

[00148] Following procedures analogous to those described in Example 1, steps 2-3, substituting 3-(N-phenylcarbamoylamino)-sulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title

compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{H}}$  9.11 (s, 1 H), 8.18 (s, 1 H), 7.73 (d,  $J = 9.0$  Hz, 2 H), 7.63 (d,  $J = 8.7$  Hz, 2 H), 7.56 (m, 1 H), 7.46 (d,  $J = 7.8$  Hz, 2 H), 7.30 (app. t,  $J = 7.8$  Hz, 2 H), 7.0 (m, 1 H), 2.80 (d,  $J = 13.5$  Hz, 2 H),  $^{31}\text{P}$  NMR (121 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{P}}$  17.0.

**Example 42: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(4-chlorophenyl) ester ammonium salt (61).**

**[00149]** Following procedures analogous to those described in Example 1, substituting 2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 4-chlorophenol for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.87 (dd,  $J = 1.5, 5$  Hz, 1 H, Ar-H), 7.73 (dd,  $J = 1.5, 4$  Hz, 1 H, Ar-H), 7.34-7.37 (m, 2 H, Ar-H), 7.23 (dd,  $J = 4, 5$  Hz, 1 H, Ar-H), 7.06-7.08 (m, 2 H, Ar-H), 3.26 (d,  $J = 27.0$  Hz, 2 H,  $\text{NCH}_2\text{P}$ );  $^{31}\text{P}$  NMR: (164 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.4;  $^{13}\text{C}$  NMR (100 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{C}}$  39.5 (d,  $J = 152$  Hz), 123.0 (d,  $J = 3.5$  Hz), 128.9, 129.8, 130.2, 134.2, 134.6, 138.1, 150.7.

**Example 43: [(3-(N-Phenylcarbamoylamino)benzenesulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (70)**

**[00150]** Following procedures analogous to those described in Example 1, substituting (N-phenylcarbamoylamino)sulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  8.08 (m, 1 H), 7.64-7.91 (m, 5 H), 7.49 (m, 2 H), 7.34 (m, 3 H), 7.08 (m, 1 H), 3.20 (d,  $J = 13.8$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  13.2.

**Example 44: [(5-Bromo-2-thiophenesulfonylamino)methyl]-phosphonic acid mono-(5-chloropyridin-3-yl) ester sodium salt (71).**

**[00151]** Following procedures analogous to those described in Example 1, substituting 5-bromo-2-thiophenesulfonyl chloride for 4-fluorobenzenesulfonyl chloride in step 2 and 5-chloro-3-hydroxypyridine for 2-chloro-3-hydroxypyridine in step 4, the title compound was obtained:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.20 (br.s. 1 H), 8.11 (br.s. 1 H), 7.54 (m, 1 H), 7.33 (d,  $J = 4.2$  Hz, 1 H), 7.05 (d,  $J = 4.2$  Hz, 1 H), 3.20 (d,  $J = 12.6$  Hz, 2 H);  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  13.9.


**Example 45: Phenylsulfonylhydrazinyl-mono-(4-nitrophenyl)-phosphonamide (189).**

**[00152]** Benzenesulfonyl hydrazide (0.5 g, 2.9 mmol) and pyridine (0.27 g, 3.2 mmol) in 50 mL of dry THF were added to a stirring solution of *p*-nitrophenylphosphoryl dichloride (0.75 g, 2.9 mmol) in



50 mL of dry THF at 0 °C. After addition was complete, the solution was brought to room temperature and stirring was continued for 4 more hours. Finally, 1 N NaOH (9 mL) was added with stirring. Volatiles were removed, and the remaining aqueous solution was freeze-dried. The resultant crude product was more than 85% pure, but final purification was accomplished by preparative HPLC on an SMT C18 column using water/acetonitrile gradient. Analytical HPLC under similar conditions showed the product to be one peak (>95% pure).  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta_{\text{H}}$  8.1 (d, 2 H, *p*-nitrophenyl), 7.4-7.65 (m, 5 H, Ar-H), 7.1 (d, 2 H, *p*-nitrophenyl);  $^{31}\text{P}$  NMR (162 MHz)  $\delta_{\text{P}}$  0.08 (s).

**Example 46: [(2-thiophenesulfonylamino)-methyl]-phosphonic acid mono-(3-chlorophenyl) ester ammonium salt (48).**



**Step 1: 2-Thiophenesulfonamide (vi).**

**[00153]** At 0 °C, a concentrated solution of ammonium hydroxide (50 mL, 1.35 mol) was added to a solution of 2-thiophenesulfonyl chloride (50 g, 0.27 mol) in methanol (300 mL). The mixture was stirred at room temperature overnight, cooled in an ice bath, and a concentrated solution of HCl was added until the pH reached 7. The resultant solution was extracted with ethyl acetate, and the combined organic layers were dried over anhydrous magnesium sulfate and concentrated to afford

45 g of a brownish solid. Trituration with methylene chloride-hexanes (1:1) afforded the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.73 (m, 1 H), 7.59 (m, 1 H), 7.05 (m, 1 H).

Step 2: [(2-Thiophenesulfonylamino)methyl]phosphonic acid dimethyl ester (vii).

**[00154]** To a solution of 2-thiophenesulfonamide (vi) (1.0 g, 6.1 mmol) in methanol (5 mL) was added a catalytic amount of sodium methoxide and paraformaldehyde (200 mg, 6.7 mmol). The reaction was heated to 55 °C for 1 h, and then the homogeneous solution was cooled and trimethyl phosphite (0.8 mL, 6.8 mmol) was added. After heating the reaction mixture for an additional 1 h at 55 °C, the reaction mixture was cooled and concentrated. The residue was dissolved in dichloromethane (15 mL), washed with water (10 mL), dried ( $\text{MgSO}_4$ ), filtered and concentrated. Purification by flash chromatography (elution with methanol-chloroform; 1:19) yielded (vii) (600 mg, 34%) as a colorless solid:  $^{31}\text{P}$  NMR: (162 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{P}}$  23.7. Also isolated was the corresponding *N*-methylated compound, [(2-thiophenesulfonyl-*N*-methylamino)methyl]-phosphonic acid dimethyl ester (234 mg, 13%) as a colorless solid:  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{C}}$  36.7, 44.6 (d,  $J = 165$  Hz), 53.2 (d,  $J = 6.5$  Hz), 127.6, 132.4, 132.6, 135.7.

Step 3: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid (43).

**[00155]** The dimethyl ester (vii) was treated with bromotrimethylsilane according to the procedure described in Example 1, step 3, to afford the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.79 (dd,  $J = 4.8, 1.2$  Hz, 1 H), 7.64 (dd,  $J = 3.9, 1.2$  Hz, 1 H), 7.17 (dd,  $J = 4.8, 3.9$  Hz, 1 H), 3.16 (d,  $J = 13.2$  Hz, 2 H).

Step 4: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid di-(3-chlorophenyl) ester (viii).

**[00156]** To a solution of 3-chlorophenol (700  $\mu\text{L}$ , 6.6 mmol) in pyridine (8 mL) at -45 °C was added thionyl chloride (230  $\mu\text{L}$ , 3.2 mmol). The reaction was stirred at -45 °C for 1 h, and then a solution of phosphonic acid (200 mg, 0.78 mmol) in pyridine (5 mL) was slowly added. The reaction mixture was warmed to ambient temperature overnight, and the solution was then concentrated. The residue was dissolved in dichloromethane (15 mL) and washed with water (10 mL). The organic extracts were dried ( $\text{MgSO}_4$ ) and concentrated, and the residue was purified by flash chromatography (elution with ethyl acetate-hexanes; 1:1) to yield the diester (viii) (86 mg, 23%) as a colorless solid:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  7.57-7.58 (m, 2 H, Ar-H), 7.14-7.26 (m, 6 H, Ar-H), 7.04-7.09 (m, 3 H, Ar-H), 6.46 (dt,  $J = 2.5, 6.5$  Hz, 1 H, NH), 3.69 (dd,  $J = 6.5, 12.5$  Hz, 2 H,  $\text{NCH}_2\text{P}$ ).

Step 5: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(3-chlorophenyl) ester ammonium salt (48).

**[00157]** To a solution of diester (viii) (86 mg, 0.18 mmol) in dioxane (1.5 mL) was added 1 *N* sodium hydroxide solution (720  $\mu$ L, 0.72 mmol) and water (4 mL). After 3 h, the solution was neutralized and concentrated. the residue was purified by flash chromatography (elution with chloroform:methanol:concentrated ammonium hydroxide; 8:2:0.25) to afford the title compound (28 mg, 40%) as a colorless solid:  $^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.84 (d,  $J$  = 5 Hz, 1 H, Ar-H), 7.71 (d,  $J$  = 3.5 Hz, Ar-H), 7.30 (t,  $J$  = 8 Hz, 1 H, Ar-H), 7.15-7.21 (m, 3 H, Ar-H), 7.01-7.03 (m, 1 H, Ar-H), 3.23 (d,  $J$  = 14 Hz, 2 H,  $\text{NCH}_2\text{P}$ );  $^{31}\text{P}$  NMR: (164 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.5.

**Example 47: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(3-dimethylamino)phenyl) ester ammonium salt (78).**

**[00158]** According to the procedure described in Example 1, step 4, the phosphonic acid **43** was treated with 3-(dimethylamino)phenol. The ammonium salt was purified by flash chromatography (elution with chloroform:methanol:concentrated ammonium hydroxide; 8:2:0.25), followed by lyophilization, to afford the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.67 (m, 1 H), 7.54 (m, 1 H), 7.11 (t,  $J$  = 7.0 Hz, 1 H), 7.04 (m, 1 H), 6.68 (d,  $J$  = 8.4 Hz, 1 H), 6.64 (s, 1 H), 6.48 (d,  $J$  = 8.4 Hz, 1 H), 3.07, d,  $J$  = 13.2 Hz, 2 H), 2.73 (s, 6 H);  $^{31}\text{P}$  NMR: (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  14.6;  $^{13}\text{C}$  NMR (75 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{C}}$  147.69, 144.91, 132.66, 129.13, 128.76, 125.97, 123.49, 110.35, 108.23, 104.94, 38.00, 34.21, (d,  $J$  = 150.0 Hz).

**Example 48: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(benzyl) ester ammonium salt (79).**

**[00159]** According to the procedure described in Example 1, step 4, the phosphonic acid **43** was treated with benzyl alcohol. The ammonium salt was purified by flash chromatography (elution with chloroform:methanol:concentrated ammonium hydroxide; 8:2:0.25), followed by lyophilization, to afford the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  7.71 (m, 1 H), 7.58 (m, 1 H), 7.29 (s, 5 H), 7.07 (t,  $J$  = 7.0 Hz, 1 H), 4.75 (d,  $J$  = 7.8 Hz, 2 H), 2.93 (d,  $J$  = 12.9 Hz, 2 H);  $^{31}\text{P}$  NMR: (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  16.65;  $^{13}\text{C}$  NMR (75 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{C}}$  132.75; 131.10, 129.07, 128.72, 124.12, 123.70, 123.47, 123.13, 62.55, 34.40, (d,  $J$  = 147.4 Hz).

**Example 49: [(2-Thiophenesulfonylamino)methyl]-phosphonic acid mono-methyl ester sodium salt (80).**

**[00160]** The diester (vii) was treated with sodium hydroxide according to the procedure described in Example 46, step 5, to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.65 (dd,  $J = 4.8, 1.2$  Hz, 1 H), 7.62 (dd,  $J = 3.6, 1.2$  Hz, 1 H), 7.15 (dd,  $J = 4.8, 3.6$  Hz, 1 H), 3.55 (d,  $J = 9.3$  Hz, 3 H), 3.04 (d,  $J = 13.8$  Hz, 2 H);  $^{31}\text{P}$  NMR: (121 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  17.60;  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  141.77, 132.98, 132.90, 128.43, 52.02, 40.11, (d,  $J = 147.2$  Hz).

**Example 50: [(3-thiophenesulfonylamino)methyl]-phosphonic acid (ix).**

**[00161]** Starting with 3-thiophenesulfonyl chloride, procedures analogous to those described in Example 46, steps 1-3, were followed to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  8.15 (dd,  $J = 3, 1.2$  Hz, 1 H), 7.66 (dd,  $J = 5.4, 3.0$  Hz, 1 H), 7.44 (dd,  $J = 5.4, 1.2$  Hz, 1 H), 3.18 (d,  $J = 13.5$  Hz, 2 H);  $^{13}\text{C}$  NMR (75.4 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  140.8, 131.8, 129.4, 126.6, 40.7 (d,  $J = 157$  Hz).

**Example 51: [(3-Thiophenesulfonylamino)methyl]-phosphonic acid mono(3-pyridyl) ester sodium salt (55).**

**[00162]** Phosphonic acid compound (ix) was treated with 3-hydroxypyridine according to the procedure described in Example 1, step 4, to provide the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.15 (br s, 2 H), 8.03 (m, 1 H), 7.49 (m, 1 H), 7.44 (br d,  $J = 8.7$  Hz, 1 H), 7.28 (m, 1 H), 7.25 (m, 1 H), 3.10 (d,  $J = 13.0$  Hz, 2 H).

**Example 52: [(3-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(2-chloro-3-pyridyl) ester sodium salt (56).**

**[00163]** Phosphonic acid compound (ix) was treated with 2-chloro-3-hydroxypyridine according to the procedure described in Example 1, step 4, to provide the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.05 (dd,  $J = 3.0, 1.2$  Hz, 1 H), 7.98 (br d,  $J = 4.8$  Hz, 1 H), 7.57 (m, 1 H), 7.48 (dd,  $J = 5.4, 3.0$  Hz, 1 H), 7.25 (m, 2 H), 3.18 (d,  $J = 12.6$  Hz, 2 H).

**Example 53: [(3-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (57).**

**[00164]** Phosphonic acid compound (ix) was treated with 4-nitrophenol according to the procedure described in Example 1, step 4, to provide the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.10 (d,  $J = 9.3$  Hz, 2 H), 8.04 (m, 1 H), 7.50 (dd,  $J = 5.1, 3.0$  Hz, 1 H), 7.26 (dd,  $J = 5.1, 1.5$  Hz, 1 H), 7.14 (d,  $J = 9.3$  Hz, 2 H), 3.11 (d,  $J = 12.9$  Hz, 2 H).

**Example 54: [(3-Thiophenesulfonylamino)methyl]-phosphonic acid mono-(2-fluoro-4-nitrophenyl) ester sodium salt (69).**

**[00165]** Phosphonic acid compound (ix) was treated with 2-fluoro-4-nitrophenol according to the procedure described in Example 1, step 4, to provide the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.05 (m, 1 H), 7.99 (m, 1 H), 7.93 (m, 1 H), 7.49 (m, 1 H), 7.27 (m, 2 H), 3.18 (d,  $J = 12.6$  Hz, 2 H).

**Example 55: [(3-Furysulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (74).**

Step 1: 3-Furysulfonamide (x)

**[00166]** 3-Furysulfonyl chloride was treated with ammonia according to the procedure described in Example 46, step 1, to produce the title compound.

Step 2: [(3-Furysulfonylamino)methyl] phosphonic acid (73).

**[00167]** 3-Furysulfonamide (x) was treated with paraformaldehyde and trimethylphosphite, followed by treatment with bromotrimethylsilane, according to the procedure described in Example 46, steps 2 and 3, to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  8.13 (dd,  $J = 1.5, 0.6$  Hz, 1 H), 7.71 (t,  $J = 1.8$  Hz, 1 H), 6.78 (dd,  $J = 1.8, 0.6$  Hz, 1 H), 3.21 (d,  $J = 13.5$  Hz, 2 H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  147.2, 146.4, 128.0, 109.3, 40.7 (d,  $J = 158$  Hz).

Step 3: [(3-Furysulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester sodium salt (74).

**[00168]** Compound 73 was treated with 4-nitrophenol in the presence of  $\text{CCl}_3\text{CN}$  according to the procedure described in Example 1, step 4, to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.11 (d,  $J = 9.0$  Hz, 2 H), 7.80 (m, 1 H), 7.51 (m, 1 H), 7.16 (d,  $J = 9.0$  Hz, 2 H), 6.61 (m, 1 H), 3.16 (d,  $J = 13.0$  Hz, 2 H).

**Example 56: [(3-Furylsulfonylamino)methyl]-phosphonic acid mono-(2-fluoro-4-nitrophenyl) ester sodium salt (75).**

[00169] Compound **73** was treated with 2-fluoro-4-nitrophenol in the presence of  $\text{CCl}_3\text{CN}$  according to the procedure described in Example 1, step 4, to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.00 (m, 3 H), 7.53 (m, 1 H), 7.35 (t,  $J = 8.4$  Hz, 1 H), 6.63 (m, 1 H), 3.21 (d,  $J = 12.9$  Hz, 2 H).

**Example 57: [(3-Furylsulfonylamino)methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester sodium salt (76).**

[00170] Compound **73** was treated with 2-chloro-3-hydroxypyridine in the presence of  $\text{CCl}_3\text{CN}$  according to the procedure described in Example 1, step 4, to produce the title compound:  $^1\text{H}$  NMR (300 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{H}}$  8.01 (m, 1 H), 7.99 (m, 1 H), 7.62 (dt,  $J = 8.1, 1.5$  Hz, 1 H), 7.51 (t,  $J = 2.0$  Hz, 1 H), 7.27 (dd,  $J = 8.1, 4.8$  Hz, 1 H), 6.61 (dd,  $J = 2.0, 0.9$  Hz, 1 H), 3.20 (d,  $J = 12.6$  Hz, 2 H).

**Example 58: [(2-benzothiophenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (77).**

Step 1: 2-Benzothiophenesulfonamide

[00171] Preparation of 2-benzothiophenesulfonamide was effected using the reported procedure (S. L. Graham et al., *J. Med. Chem.* **1989**, 32, 2548).

Step 2: [(2-Benzothiophenesulfonylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (77).

[00172] 2-Benzothiophenesulfonamide was treated with paraformaldehyde and trimethylphosphite, followed by treatment with bromotrimethylsilane, according to the procedure described in Example 46, steps 2 and 3. The crude product was purified by flash chromatography (elution with dichloromethane:methanol: concentrated ammonium hydroxide; 7:3.5:0.5), followed by lyophilization, to afford the ammonium salt **77**:  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  7.9 (m, 5 H, Ar-H), 7.4 (m, 2 H, Ar-H), 7.2 (d,  $J = 9$  Hz, 2 H, Ar-H), 2.8 (d,  $J = 12.9$  Hz, 2 H,  $\text{CH}_2$ );  $^{31}\text{P}$  NMR (121 MHz,  $\text{D}_2\text{O}$ )  $\delta_{\text{P}}$  12.08 (t,  $J = 12.2$  Hz, 1 P).

**Example 59: [(2-Thiophenesulfonyl-N-methylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (47).**


Step 1: [(2-Thiophenesulfonyl-N-methylamino)methyl]-phosphonic acid (xi).

**[00173]** To a solution of [(2-thiophenesulfonyl-N-methylamino)methyl]-phosphonic acid dimethyl ester (obtained as described in Example 46, step 2) (180 mg, 0.6 mmol) in dichloromethane (3 mL) at 0 °C was slowly added bromotrimethylsilane (190  $\mu$ L, 1.4 mmol). The reaction was gradually warmed to room temperature and allowed to stir overnight. The reaction mixture was concentrated, the residue was redissolved in dichloromethane, and the solution was again concentrated. Methanol (10 mL) was added to the residue, and the solution was stirred at room temperature for 30 min. The solution was then filtered and concentrated. The oily residue was triturated with 1:1 ether-hexanes, and an orange solid was collected by filtration. The orange solid was boiled in dichloromethane (3 mL), cooled, and filtered to yield the title compound (120 mg, 74%) as a colorless solid:  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  7.87 (dd,  $J = 1.5, 4$  Hz, 1 H, Ar-H), 7.66 (dd,  $J = 1.5, 3.5$  Hz, 1 H, Ar-H), 7.25 (dd,  $J = 4, 5$  Hz, 1 H, Ar-H), 3.31 (d,  $J = 12.5$  Hz, 2 H,  $\text{CH}_2\text{P}$ ), 2.92 (s, 3 H,  $\text{NCH}_3$ ).

Step 2: [(2-Thiophenesulfonyl-N-methylamino)methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (47).

**[00174]** Compound (xi) was treated with 4-nitrophenol according to the procedure described in Example 1, step 4. The crude product was purified by flash chromatography (elution with chloroform:methanol:concentrated ammonium hydroxide; 8:2:0.25) to afford the ammonium salt **47** (43%) as a colorless solid:  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  8.19-8.23 (m, 2 H, Ar-H), 7.85 (dd,  $J = 1, 5$  Hz, 1 H, Ar-H), 7.62 (dd,  $J = 1, 4$  Hz, 1 H, Ar-H), 7.44 (dd,  $J = 1, 9.5$  Hz, 2 H, Ar-H), 7.23 (dd,  $J = 4, 5$  Hz, 1 H, Ar-H), 3.30 (d,  $J = 13.5$  Hz, 2 H,  $\text{NCH}_2\text{P}$ ), 2.93 (s, 3 H,  $\text{NCH}_3$ );  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  37.1, 47.4, 122.4 (d,  $J = 4$  Hz), 126.1, 128.9, 133.7 (d,  $J = 7$  Hz), 137.1, 144.7, 159.5;  $^{31}\text{P}$  NMR (162 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  12.7; MS (neg. FAB)  $m/z$  391 (M - 1, 100%), 306 (15%), 153 (35%).

**Example 60: [2-Phenyl-1-(thiophene-2-sulfonylamino)-ethyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt. (44)**



**Step 1: [2-Phenyl-1-(thiophene-2-sulfonylamino)-ethyl]-phosphonic acid diethyl ester (xii).**

**[00175]** To a suspension of sulfonamide (vi) (500 mg; 0.31 mmol) in acetyl chloride (5 mL) was slowly added diethyl phosphite (400  $\mu$ L; 0.31 mmol). The reaction was cooled to 0°C and then phenylacetaldehyde (450  $\mu$ L; 0.39 mmol) was added dropwise. The reaction was gradually warmed to ambient temperature and after 6 hrs the homogeneous solution was concentrated. The residue was diluted with dichloromethane (25 mL) then washed successively with water (10 mL), saturated sodium bicarbonate (10 mL) and brine (10 mL). The combined organic extracts were dried (MgSO<sub>4</sub>), evaporated and initially purified by flash chromatography (ethyl acetate-hexanes; 3:2). A second purification by flash chromatography (acetone-hexanes; 3:7) yielded diethyl phosphonate (xii) (437 mg, 35%) as a colorless solid; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>H</sub> 1.22 (t, J = 7 Hz, 3 H, OCH<sub>2</sub>CH<sub>3</sub>), 1.26 (t, J = 7 Hz, 3 H, OCH<sub>2</sub>CH<sub>3</sub>), 2.89 (dt, J = 8, 14 Hz, 1 H, CHPh), 3.12 (dt, J = 6, 14 Hz, 1 H, CHPh), 3.96-4.23 (m, 5 H, NCHP and OCH<sub>2</sub>CH<sub>3</sub>), 6.38 (dd, J = 2, 9.5 Hz, 1 H, NH), 6.92-6.93 (m, 1 H, Ar-H), 7.12-7.20 (m, 5 H, Ar-H), 7.33-7.34 (m, 1 H, Ar-H), 7.43-7.45 (m, 1 H, Ar-H).

**Step 2: [2-Phenyl-1-(thiophene-2-sulfonylamino)-ethyl]-phosphonic acid (xiii).**

**[00176]** To a solution of diethyl phosphonate (xii) (745 mg; 1.8 mmol) in dichloromethane (8 mL) at 0 °C was slowly added bromotrimethylsilane (2 mL; 15.2 mmol). The reaction was gradually warmed to room temperature and the following day the solution was evaporated. The residue was redissolved in dichloromethane (10 mL), evaporated and the process repeated once more. Methanol (10 mL) was added to the residue and the solution stirred at room temperature for 30 minutes, then



the solution was filtered and concentrated. The oily residue was triturated with 1:1 ether-hexanes and the orange solid collected by filtration. Finally, the orange solid was boiled in dichloromethane (5 mL), cooled and filtered to yield pure phosphonic acid (**xiii**) (512 mg, 80%) as a colorless solid:  $^1\text{H}$  NMR (400MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  2.71 (dt,  $J = 14, 10$  Hz, 1 H,  $\text{CHPh}$ ), 3.13 (ddd,  $J = 4, 8, 14$  Hz, 1 H,  $\text{CHPh}$ ), 3.89 (ddd,  $J = 4, 10, 14$  Hz,  $\text{NCHP}$ ), 6.86 (dd,  $J = 3.5, 5$  Hz, 1 H, Ar-H), 7.07-7.15 (m, 6 H, Ar-H), 7.53 (dd,  $J = 1.5, 5$  Hz, 1 H, Ar-H).

**Step 3: [2-Phenyl-1-thiophene-2-sulfonylamino)-ethyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (**44**)**

**[00177]** To sealed tube containing phosphonic acid (**xiii**) (50 mg; 0.14 mmol) and sublimed 4-nitrophenol (24 mg; 0.17 mmol) was added pyridine (0.7 mL) and trichloroacetonitrile (100  $\mu\text{L}$ ; 1.0 mmol). The reaction was heated to 105  $^{\circ}\text{C}$  for 6 hrs then was cooled and concentrated. The residue was purified by flash chromatography (chloroform-methanol-concentrated ammonium hydroxide; 8:2:0.25) to yield the ammonium salt **44** (36.4 mg, 52%) as a colorless solid:  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  2.83 (td,  $J = 9, 14$  Hz, 1 H,  $\text{CHPh}$ ), 3.23-3.29 (m, 1 H,  $\text{CHPh}$ ), 3.95-4.02 (m, 1 H,  $\text{NCHP}$ ), 6.85 (dd,  $J = 4, 5$  Hz, 1 H, Ar-H), 7.08-7.20 (m, 6 H, Ar-H), 7.27-7.31 (m, 2 H, Ar-H), 7.52 (dd,  $J = 1.5, 5$  Hz, 1 H, Ar-H), 8.12-8.16 (m, 2 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  16.7;  $^{13}\text{C}$  NMR: (100MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{C}}$  38.1, 122.0, 125.9, 127.3, 128.3, 129.2, 130.6, 132.2, 132.3, 144.4, 144.5.

**Example 61: [2-Methyl-1-thiophene-2-sulfonylamino)-propyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (**52**)**

**[00178]** The title compound was synthesized according to the three-step procedure described in Example 60, Steps 1-3, but substituting isobutyraldehyde for phenylacetaldehyde. The mono ester **52** was isolated as a colorless solid (13% yield over three steps):  $^1\text{H}$  NMR: (400 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  0.86 (d,  $J = 7$  Hz, 3 H,  $\text{CH}_3$ ), 0.93 (d,  $J = 7$  Hz, 3 H,  $\text{CH}_3$ ), 2.02-2.10 (m, 1 H,  $(\text{CH}_3)_2\text{CH}$ ), 3.28-3.35 (m, 1 H,  $\text{NCHP}$ ), 7.03 (dd,  $J = 3.5, 5$  Hz, 1 H, Ar-H), 7.12 (br s, 5 H, NH), 7.17 (d,  $J = 9$  Hz, 2 H, Ar-H), 7.60 (dd,  $J = 1, 3.5$  Hz, 1 H, Ar-H), 7.75 (dd,  $J = 1, 5$  Hz, 1 H, Ar-H), 8.06-8.10 (m, 2 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{P}}$  14.2;  $^{13}\text{C}$  NMR: (100MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{C}}$  19.8, 21.3, (d,  $J = 10$ ), 30.7, 58.3 (d,  $J = 142.3$ ), 121.3 (d,  $J = 4.5$ ), 125.9, 128.2, 132.2, 132.4, 142.2, 144.7, 161.2.

**Example 62: [Phenyl-(thiophene-2-sulfonylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (53).**

**[00179]** The title compound was synthesized according to the three-step procedure in Example 60, Steps 1-3, but substituting benzaldehyde for phenylacetaldehyde. The mono ester **53** was isolated as a colorless solid (8% yield over three steps):  $^1\text{H}$  NMR (400 MHz, DMSO- $\text{D}_6$ )  $\delta_{\text{H}}$  4.44 (d, J = 22 Hz, 1 H, NCHP), 6.83 (dd, J = 4, 5 Hz, 1 H, Ar-H), 7.00-7.07 (m, 4 H), 7.17-7.33 (m, 9 H), 7.60 (dd, J = 1.5, 5 Hz, 1 H, Ar-H), 8.08-8.12 (m, 2 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz, DMSO- $\text{D}_6$ )  $\delta_{\text{P}}$  10.1;  $^{13}\text{C}$  NMR: (100MHz, DMSO- $\text{D}_6$ )  $\delta_{\text{C}}$  58.0 (d, J = 140.2 Hz) 121.6 (d, J = 4 Hz), 125.9, 126.9, 127.9, 128.1, 129.1 (d, J = 5 Hz), 132.3, 132.6, 138.9, 142.4, 143.6 and 161.4.

**Example 63: [2-Phenyl-1-(thiophene-2-sulfonylamino)-ethyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester ammonium salt (46).**

**[00180]** To a sealed tube containing phosphonic acid (**xiii**) (134 mg; 0.39 mmol) and 2-chloro-3-pyridinol (60 mg; 0.46 mmol) was added pyridine (1.9 mL) and trichloroacetonitrile (270  $\mu\text{L}$ ; 2.7 mmol). The reaction was heated to 105  $^{\circ}\text{C}$  for 6 hours then was cooled and concentrated. The residue was purified by flash chromatography (chloroform-methanol-concentrated ammonium hydroxide; 9:2:0.1) to yield the ammonium salt **46** (100 mg, 54%) as a colorless solid:  $^1\text{H}$  NMR: (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  2.87 (td, J = 9, 14 Hz, 1 H, CHPh), 3.28-3.36 (m, 1 H, CHPh), 4.01 (ddd, J = 4.5, 9, 17 Hz, 1 H, NCHP), 6.81 (dd, J = 3.5, 5 Hz, 1 H, Ar-H), 7.07-7.21 (m, 6 H, Ar-H), 7.28 (dd, J = 4.5, 8 Hz, 1 H, Ar-H), 7.46 (dd, J = 1.5, 5 Hz, 1 H, Ar-H), 7.91-7.93 (m, 1 H, Ar-H), 7.99-8.02 (m, 1 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{P}}$  17.0;  $^{13}\text{C}$  NMR: (100MHz, DMSO- $\text{D}_6$ )  $\delta_{\text{C}}$  38.3, 55.6, (d, J = 147.6 Hz), 124.3, 126.7, 128.1, 130.1, 130.4, 131.8, 132.1, 140.3 (d, J = 10 Hz), 142.4, 142.7, 144.5, 147.9.

**Example 64: [2-Methyl-1-(thiophene-2-sulfonylamino)-propyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester ammonium salt (54)**


**[00181]** Following the same procedure as described in Example 63, but substituting the phosphonic acid derived from Example 61 for phosphonic acid (**xiii**), the title compound was obtained. The mono ester **54** was isolated as a colorless solid (37% yield):  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  1.02 (d, J = 7 Hz, 3 H,  $\text{CH}_3$ ), 1.06 (d, J = 7 Hz, 3 H,  $\text{CH}_3$ ), 2.24-2.32 (m, 1 H,  $(\text{CH}_3)_2\text{CH}$ ), 3.69 (dd, J = 3.5, 19.5 Hz, 1 H, NCHP), 6.91 (dd, J = 4, 5 Hz, 1 H, Ar-H), 7.25 (dd, J = 5, 8 Hz, 1

H, Ar-H), 7.52-7.59 (m, 2 H, Ar-H), 7.80-7.82 (m, 1 H, Ar-H), 7.99-8.01 (m, 1 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{p}}$  17.3;  $^{13}\text{C}$  NMR: (100 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{c}}$  18.8, 21.1 (d,  $J = 11.5$ ), 31.1 (d,  $J = 5$ ), 59.0 (d,  $J = 149.5$ ), 124.5, 128.0, 131.2, 132.2, 132.8, 143.6, 144.5, 156.2, 156.4.

**Example 65: [Phenyl-(thiophene-2-sulfonylamino)-methyl]-phosphonic acid mono-(2-chloropyridin-3-yl) ester ammonium salt (60)**

**[00182]** Following the same procedure as described in Example 63, but substituting the phosphonic acid derived from Example 62 for phosphonic acid (xiii), the title compound was obtained. The mono ester **60** was isolated as a colorless solid (50% yield):  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  4.49 (d,  $J = 22$  Hz, 1 H, NCHP), 6.80 (dd,  $J = 4, 5$  Hz, 1 H, Ar-H), 6.99-7.07 (m, 3 H, Ar-H), 7.20-7.33 (m, 9 H), 7.58 (dd,  $J = 1.5, 5$  Hz, 1 H, Ar-H), 7.84 (dd,  $J = 1.5, 8$  Hz, 1 H, Ar-H), 7.94 (dd,  $J = 1.5, 5$  Hz, 1 H, Ar-H);  $^{31}\text{P}$  NMR: (162MHz,  $\text{DMSO}-D_6$ )  $\delta_{\text{p}}$  10.7;  $^{13}\text{C}$  NMR: (100 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{c}}$  58.3 (d,  $J = 140.5$ ), 124.3, 127.0, 127.9, 128.1, 129.1 (d,  $J = 5$ ), 129.9, 132.3, 132.6, 138.8, 142.5, 142.9, 143.5, 148.0.

**Example 66: [(2-Benzothiophenesulfonyl-N-methylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt. (142)**



Step 1: [(2-Benzothiophenesulfonyl-N-methylamino)-methyl]-phosphonic acid dimethyl ester (xiv).

**[00183]** A reaction mixture consisting of 290 mg (0.83 mmol) of [(2-benzothiophenesulfonylamino)-methyl]-phosphonic acid dimethyl ester, 380 mg (1.16 mmol) of anhydrous cesium carbonate, 0.11 mL (1.16 mmol) of methyl iodide and 5 mL of anhydrous DMF stirred for 16 hours at room temperature. After the reaction was complete, DMF was evaporated in vacuum, and the residue was suspended in 15 mL of water and extracted with dichloromethane. The

extract was dried over anhydrous magnesium sulfate, filtered and evaporated, providing an oily residue which was chromatographed on a silica gel column, eluent EtOAc/hexane (7:3), then dichloromethane/MeOH (20:1), to yield the title compound as a yellowish oil (190 mg, 63%):  $^1\text{H}$  NMR: (300 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  3.00 (s,  $\text{CH}_3$ ), 3.48 (d,  $J = 11.5$  Hz, 2 H), 3.83 (d,  $J = 10.9$  Hz, 6 H,  $2\text{CH}_3$ ), 7.45-7.50 (m, 2 H, Ar-H), 7.83-7.89 (m, 3 H, Ar-H).

**Step 2: [(2-Benzothiophenesulfonyl-N-methylamino)-methyl]-phosphonic acid (xv)**

**[00184]** To a solution of 550 mg (1.58 mmol) of the compound (xiv) in 10 mL of anhydrous dichloromethane at 0 °C, 0.49 mL of bromotrimethylsilane was added. The mixture was warmed up to room temperature and left overnight. Dichloromethane was evaporated; the oily residue was dissolved in another 10 mL of dichloromethane, filtered, again evaporated and redissolved in 10 mL of MeOH. The methanolic solution stirred at room temperature 30 min., evaporated and the residue was triturated with a mixture ether/hexane to form a white crystalline product which was separated by filtration (390 mg, 70%):  $^1\text{H}$  NMR: (300 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  2.88 (s,  $\text{CH}_3$ ), 3.21 (d,  $J = 12.1$  Hz, 2 H), 7.49-7.61 (m, 2 H, Ar-H), 8.01-8.17 (m, 3 H, Ar-H).

**Step 3: [(2-Benzothiophenesulfonyl-N-methylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (142)**

**[00185]** A reaction mixture consisting of 500 mg (1.56 mmol) of (xv), 265mg (1.90 mmol) of p-nitrophenol, 1.1 mL (11 mmol) of trichloroacetonitrile, and 5 mL of anhydrous pyridine was stirred in a sealed tube for 5 hours at 105-110 °C. After the reaction was complete, pyridine was evaporated in vacuum, the oily residue was chromatographed on a silica gel column, eluent chloroform/MeOH/ammonium hydroxide (9:2:0.1) to yield 620 mg (87%) of the title compound as a pale crystalline substance:  $^1\text{H}$  NMR: (300 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  2.89 (s,  $\text{CH}_3$ ), 3.07 (d,  $J = 11.5$  Hz, 2 H), 7.15 (br s, 4 H,  $\text{NH}_4$ ), 7.37 (d,  $J = 9.3$  Hz, 2 H, Ar-H), 7.47-7.57 (m, 2 H, Ar-H), 8.01-8.14 (m, 5 H, Ar-H).

**Example 67: [(2-Benzothiophenesulfonyl-N-ethylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (144)**

**[00186]** The title compound was prepared in 62% yield following the procedure from Example 66, steps 1-3, but using ethyl iodide as the alkylating agent:  $^1\text{H}$  NMR: (300 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  1.00 (t,  $J = 7.0$  Hz,  $\text{CH}_3$ ), 3.33 (d,  $J = 11.1$  Hz,  $\text{PCH}_2$ ), 3.49 (q,  $J = 7.1$  Hz), 6.95-7.51 (m, 8 H, Ar-H,  $\text{NH}_4$ ), 7.98-8.10 (m, 5 H, Ar-H).


**Example 68: [(2-Benzothiophenesulfonyl-N-benzylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (145).**

**[00187]** The title compound was prepared in 58% yield following the procedure from Example 66, steps 1-3 but using benzyl bromide as the alkylating agent:  $^1\text{H}$  NMR: (300 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  3.29 (d,  $J = 10.0$  Hz,  $\text{PCH}_2$ ), 4.78 (s,  $\text{CH}_2\text{Ph}$ ), 7.16-7.47 (m, 13 H, Ar-H,  $\text{NH}_4$ ), 7.96-8.00 (m, 4 H, Ar-H), 8.07 (s, 1 H, Ar-H)

**Example 69: [(2-Benzothiophenesulfonyl-N-phenylethylamino)-methyl]-phosphonic acid mono-(4-nitrophenyl) ester ammonium salt (143)**

**[00188]** The title compound was prepared in 55% following the procedure from Example 66, steps 1-3, but using 2-bromoethyl benzene as the alkylating agent:  $^1\text{H}$  NMR: (300 MHz, DMSO- $d_6$ )  $\delta_{\text{H}}$  2.86 (dd,  $J = 8.4$ , Hz,  $\text{NCH}_2$ ), 3.43 (d,  $J = 11.2$  Hz,  $\text{PCH}_2$ ), 3.58 (dd,  $J = 8.5$  Hz,  $\text{CH}_2\text{Ph}$ ), 7.13-7.56 (m, 13 H, Ar-H,  $\text{NH}_4$ ), 7.96-8.10 (m, 5 H, Ar-H).

**Example 70: (5,8-dichlorobenzothiophene-2-sulfonyl)aminomethylphosphonic acid mono-(4-nitrophenyl) ester ammonium salt (128).**



**Step 1: 2,2-Diethoxyethyl 2,5-dichlorophenyl sulfide (xvi).**

**[00189]** Under a nitrogen atmosphere, to 2,5-dichlorobenzenethiol (25 g, 139 mmol) in 140 mL of acetone in a 250-mL round-bottomed flask was added 21 g (154 mmol) of potassium carbonate, followed by 22 mL (146 mmol) of diethyl acetal. The mixture was stirred overnight at room temperature and then filtered, washing the collected solid with acetone, and the filtrate was concentrated. Water was added to the concentrated filtrate, and the aqueous phase was extracted

three times with ethyl acetate. The combined organic layers were washed with 0.5 M KOH, water and brine, dried over magnesium sulfate, filtered, and concentrated. The title compound was obtained as a yellow oil (37 g, 90%):  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  7.36 (d,  $J = 2.1$  Hz, 1 H), 7.26 (d,  $J = 8.4$  Hz, 1 H), 7.06 (dd,  $J = 2.1, 8.4$  Hz, 1 H), 4.71 (t,  $J = 5.4$  Hz, 1 H), 3.71 (quintuplet,  $J = 7.2$  Hz, 2 H), 3.57 (quintuplet,  $J = 6.9$  Hz, 2 H), 3.14 (d,  $J = 5.4$  Hz, 2 H), 1.22 (t,  $J = 7.2$  Hz, 6 H).

Step 2: 5,8-Dichlorobenzothiophene (xvii).

**[00190]** Anhydrous chlorobenzene (250 mL) was placed in a 500-mL three-necked flask equipped with a condenser, an addition funnel, and a magnetic stirring bar. The apparatus was flushed with nitrogen, and polyphosphoric acid (40g) was added. The mixture was heated at 125 °C, and 2,2-diethoxyethyl 2,5-dichlorophenyl sulfide (24 g, 82 mmol) in 25 mL of chlorobenzene was added over one hour. The resultant mixture was allowed to stir overnight. The reaction mixture was cooled to room temperature, and the organic phase was separated from polyphosphoric acid. Residual polyphosphoric acid was decomposed with water, and the resultant aqueous phase was washed twice with chloroform. The combined organic phases were dried over magnesium sulfate and concentrated. The crude product was purified by flash chromatography, eluting with a mixture of 5% ethyl acetate: 95% hexane to give 1.5 g (99%) of 5,8-dichlorobenzothiophene as a light yellow solid:  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  7.57 (d,  $J = 5.4$  Hz, 1 H), 7.53 (d,  $J = 5.4$  Hz, 1 H), 7.32 (d,  $J = 8.1$  Hz, 1 H), 7.27 (d,  $J = 8.1$  Hz, 1 H).

Step 3: 5,8-Dichlorobenzothiophene-2-sulfonamide (xviii).

**[00191]** Under a nitrogen atmosphere, to 5,8-dichlorobenzothiophene (6.58 g, 32 mmol) in 50 mL of THF at -78 °C in a 250-mL round-bottomed flask was slowly added a 2.5 M solution of *n*-butyllithium in pentane (16 mL, 40 mmol). After 30 minutes, sulfur dioxide gas was bubbled over the solution until the mixture became acidic. Hexane was added, and the mixture was warmed to room temperature. The sulfinic salt was collected by filtration and dried overnight under high vacuum.

**[00192]** Under a nitrogen atmosphere, to the sulfinic salt (8.88 g, 32 mmol) in 60 mL of dichloromethane at 0 °C in a 250-mL round-bottomed flask was added *N*-chloro-succinimide (6.29 g, 47 mmol). After 15 minutes, the mixture was warmed to room temperature over 45 minutes. The suspension was filtered over Celite, and the filtrate was concentrated to give 9.65 g (100%) of the corresponding sulfonyl chloride. This material was dissolved in 50 mL of acetone and cooled to 0 °C, and 10 mL of ammonium hydroxide was added. After 15 minutes, the mixture was warmed to

room temperature over 45 minutes, and the solvents were evaporated. The residue was dissolved in water and extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated. The title compound was obtained as a white solid (6.45 g, 70%):  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  8.09 (s, 1 H), 7.41 (s, 2 H)

Step 4: (5,8-dichlorobenzothiophene-2-sulfonyl)aminomethylphosphonic acid mono-(4-nitrophenyl) ester ammonium salt (128).

[00193] Compound **128** was prepared by procedures analogous to those described in Example 46, steps 2 and 3, and Example 1, step 4:  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta_{\text{H}}$  7.94 (s, 1 H), 7.92 (d,  $J = 9.3$  Hz, 2 H), 7.57-7.65 (m, 2 H), 7.21-7.38 (m, 4 H), 7.22 (d,  $J = 9.3$  Hz, 2 H), 2.96 (d,  $J = 12.9$  Hz, 2 H).

**Example 71: Determination of  $\text{IC}_{50}$ .**

[00194] Enzyme activity against the commercially available substrate, nitrocefin, was measured in the presence of various inhibitor concentrations. The P99  $\beta$ -lactamase enzyme (1 mg/mL) was dissolved in 20 mM MOPS buffer at pH 7.50, and diluted 200-fold in the same buffer containing 0.1% BSA. Nitrocefin was stored as a 100  $\mu\text{M}$  stock solution in 20 mM MOPS at pH 7.5. Stock solutions of test compounds were prepared in 10 mM MOPS at pH 7.5 (approximately 1 mg/mL)

[00195] A typical assay contained 300  $\mu\text{L}$  of 100  $\mu\text{M}$  nitrocefin,  $x$   $\mu\text{L}$  of inhibitor test compound,  $110-x$   $\mu\text{L}$  of MOPS, and 10  $\mu\text{L}$  of P99  $\beta$ -lactamase. The reaction at 25  $^{\circ}\text{C}$  was followed spectrophotometrically at 482 nm.  $\text{IC}_{50}$  values were determined from the dose-response plots of the initial rates vs. inhibitor concentration

[00196] TEM R+ and L-1  $\beta$ -lactamases were assayed as described above for P99  $\beta$ -lactamase.

[00197] Representative results using the test inhibitors of the invention showing the inhibition of  $\beta$ -lactamases are shown in Tables 1, 2, 4, and 5.

**Example 72: Determination of Antibiotic Minimum Inhibitory Concentration (MIC) in the Presence and Absence of  $\beta$ -Lactamase Inhibitor.**

Principle

[00198] An increasing number of organisms produce enzymes that inhibit the actions of  $\beta$ -lactam antibiotics, so-called  $\beta$ -lactamases. The "strength" of  $\beta$ -lactam antibiotics can be improved by pairing them with compounds that inhibit bacterial enzyme  $\beta$ -lactamases. The Minimum Inhibitory

Concentration (MIC) test determines the lowest concentration of an antibiotic at which no visible bacterial growth occurs. This test permits comparisons to be made between bacterial growth in the presence of antibiotic alone (control) and bacterial growth in the presence of both an antibiotic and a novel (test) compound.

#### Materials required

- β-lactamase positive strains
- β-lactam antibiotics
- β-lactamase inhibitor compounds
- appropriate bacterial growth medium

#### β-Lactamase positive strains

- Klebsiella oxytoca [ATCC#51983]
- Staphylococcus aureus [MGH: MA#50848]
- Enterococcus faecalis [ATCC#49757]
- Enterobacter cloacae [ATCC#23355]
- Haemophilus influenzae [ATCC#43163]

#### β-Lactam antibiotics

Piperacillin	333 mg/mL
TAZOCIN™ (piperacillin + tazobactam)	308 mg/mL
Ticarcillin	500 mg/mL
TIMENTIN™ (ticarcillin + clavulanic acid)	500 mg/mL
Cefoxitin	333 mg/mL
Ceftriaxone	250 mg/mL
Preparation:	Dissolved in MHB (Mueller Hinh Broth)
Working dilutions:	0.06 mg/mL to 4.22 mg/mL

#### β-Lactamase inhibitor compounds (test compounds)

Preparation:	Dissolved in MHB or DMSO
Stock solution:	2 mg/mL
Working concentration:	1, 10, 100 µg/mL

#### Culture media



MHB:	Mueller Hinth Broth
HTM:	Haemophilus Test Medium Broth
origin:	Becton Dickinson
storage:	4 °C
Blood agar (prepared general culture plate)	
Chocolate agar (prepared plate + accessory factors, required by <i>H. influenzae</i> )	
origin:	Oxoid
storage:	4 °C

### Assay Procedure

**[00199]** Frozen bacterial stocks were thawed to room temperature, and a few drops were placed on an appropriate plate; chocolate agar was used for *H. influenzae*, blood agar was used for all others. The plates were incubated overnight at 37 °C in air, except for *H. influenzae*, which was incubated in a CO<sub>2</sub> incubator. The cultures were subcultured the following day, and the subcultures were again incubated overnight

**[00200]** A sterile loop was touched to 3 colonies and used to inoculate 1 mL of Mueller-Hinton Broth (MHB). The bacterial solution was diluted 18.8-fold (20 µL of bacterial solution in 355 µL of broth). 5 µL of the resultant inoculum solution contained  $\sim 4 \times 10^4$  CFU/mL. Confirmation of the inoculum was performed by preparing a serial dilution to give 100 µL of a 5000-fold dilution, all of which was then plated onto an appropriate plate. After incubation overnight at 37 °C, the plate had 75-150 colonies.

**[00201]** In each well of a 96-well microtiter plate was combined 5 µL of bacterial inoculum, 90 µL of antibiotic dilutions, and 5 µL of test compound (or media for determination of the MIC of the antibiotic in the absence of test compound). Each plate included 2 wells with no bacterial inoculum (negative control) and 2 wells with no test compound and no antibiotic (positive control). The plate was covered and incubated with gentle shaking for 16-20 hours at 37 °C in air (CO<sub>2</sub> for *H. influenzae*). Absorbance was read at 540 nm (Multiskan Titertek mcc/340). Bacterial growth was scored as follows:

invisible to the naked eye (OD < 0.05)	NEGATIVE
barely visible (OD > 0.05, but < 50% of positive control)	PLUS/MINUS
readily visible (typically OD > 0.10)	PLUS

abundant growth (OD > 0.3)

PLUS-PLUS

The minimum inhibitory concentration (MIC), defined to be the lowest dilution of antibiotic that completely inhibits growth, was determined for the antibiotic alone and for antibiotic in the presence of each test compound. In some cases, the determination of MIC in the presence of test compound was performed at more than one concentration of the test compound.

**[00202]** Representative data is presented in Tables 1 and 2. The impact of the novel  $\beta$ -lactamase inhibitors is expressed as the ratio of the MIC determined in the absence of  $\beta$ -lactamase inhibitor to the MIC determined in the presence of the specified concentration of the  $\beta$ -lactamase inhibitor test compound. A value of 1 indicates that the  $\beta$ -lactamase inhibitor had no effect on antibiotic potency, while a positive integer indicates that the  $\beta$ -lactamase inhibitor produces a synergistic effect when co-administered with an antibiotic agent, that is, a higher concentration of antibiotic is required to completely inhibit visible bacterial growth in the absence of the test compound than in its presence

**Example 73: Synergistic Effect of  $\beta$ -Lactamase Inhibitors When Tested Against Highly Resistant  $\beta$ -Lactamase Positive Bacterial Strains**

Following procedures identical to those described in Example 72,  $\beta$ -lactamase inhibitors were tested for their ability to enhance antibiotic efficacy against  $\beta$ -lactamase positive bacterial strains that are very highly resistant to  $\beta$ -lactam antibiotics. Representative results are presented in Table 3.

Highly resistant  $\beta$ -Lactamase positive strains

Enterobacter cloacae (derepressed)

Pseudomonas aeruginosa [ATCC#12470-resistant]


Stenotrophomonas maltophilia [ATCC#12968-resistant]

Pseudomonas aeruginosa [ATCC#98043010-intermediate resistance]

Stenotrophomonas maltophilia [ATCC#98043029-intermediate resistance]

**[00203]** Successful inhibition of bacterial  $\beta$ -lactamase activity in these assays is expected to be predictive of success in animals and humans. For examples of the successful clinical development of  $\beta$ -lactamase inhibitors identified by in vitro testing, see, e.g., Di Modugno *et al.*, Current Opinion in Anti-Infective Investigational Drugs 1:26-39 (1999); Moellering, J. Antimicrobial Chemotherapy 31 Suppl. A:1-8 (1993).

**Table 1:  $\beta$ -Lactamase Inhibition and Microbiological Efficacy of Sulfonamidomethylphosphonate Derivatives.**



Cpd.	R <sup>1</sup>	R <sup>3</sup>	R <sup>4</sup>	Log P	IC <sub>50</sub> ( $\mu$ M) "C" <sup>1</sup>	Synergy <sup>2</sup> (at $\mu$ g/mL)		
						Class C E.Cl <sup>3</sup> 10/1 <sup>4</sup>	Class A H.In. 10/1 <sup>5</sup>	St.A. 10
1		H	PNP <sup>6</sup>		162	1/1	(1)	(1)
2		H	PNP		16		24	2
3		H	PNP		5	2/1	(1)	(1)
4		H	PNP		2	2/1	(1)	(1)
5		H	PNP		4	2/1	(1)	(1)

<sup>1</sup> Class C  $\beta$ -lactamase [P99].















<sup>2</sup> Expressed as the ratio of the antibiotic MIC determined in the absence of  $\beta$ -lactamase inhibitor to the antibiotic MIC determined in the presence of  $\beta$ -lactamase inhibitor.
















<sup>3</sup> E.Cl = *Enterobacter cloacae* [ATCC #23355]; H.In. = *Haemophilus influenzae* [ATCC #43163]; St.A. = *Staphylococcus aureus* [MGH:MA #50848].



















<sup>4</sup> The first value provided was determined at an inhibitor concentration of 10  $\mu$ g/mL, and the second value was determined at an inhibitor concentration of 1  $\mu$ g/mL. Where only one value is provided, it was determined at an inhibitor concentration of 10  $\mu$ g/mL.























<sup>5</sup> Values within parentheses were determined at an inhibitor concentration of 100  $\mu$ g/mL.

<sup>6</sup> PNP = *p*-nitrophenol

























6		H	PNP		7	2/1	(1)	(1)
7		H	PNP		4	4/2	(16)	(2)
8		H			>100	2/1	1	(1)
9		H	PNP		5	4/2	32	2
10		H	PNP		6	2/1	16	4
11		H			>100	1/1	8	1
12		H	PNP		10	1/1	8	1
13		H	PNP		6	4/1	8	1
14		H	PNP		3	4/2	(8)	(2)
15		H	PNP		18	3/1	4	1
16		H	PNP		19	1/1	8	1
17		H	PNP		9	2/1	8	2
























18		H	PNP		4	4/1	8	1
19		H	PNP		43	4/4	(4)	(1)
20		H	F			8/8	(16)	(1)
21		H	PNP		4	2/1	(2)	(1)
22		H	PNP		7	2/1	16	4
23		H	PNP		14	1/1	2	1
24		H	PNP		119	4/1	8	1
25		H			1,200	10/2	1	2
26		H			1,200	16/2	1	1
27		H	PNP		5.0	1/1	4	2
28		H			1,800			
29		H	F		972			























30		H			622	8/2	1	1
31		H	PNP		136	8/2	2	1
32		H			639	16/2	1	1
33		H	PNP		5	8/4	4	4
34		H	OH		1600	2/2	1	1
35		H	OCH <sub>3</sub> diester		—	1/1	1	1
36		H			410	2/2	1	1
37		H			2000	4/2	1	1
38		H			362	8/4	1	1
39		H	PNP		7	1/1	8	4
40		H			356	2/1	4	2
41		H	PNP		5	2/1	8	4



















42		H			59	8/1	2	1
43		H	OH					
44			PNP		192	2/2	1	1
45		H			3500	2/1	1	1
46					188	1/1	1	1
47		H N-Me	PNP		276	2	1	
48		H			189	4/2	1	1
49		H			3000	1/1	1	1
50		H			3000	2/1	1	1
51		H	OMe		1700	2/1	1	1
52			PNP		1400	4/2	1	1
53			PNP		363	2/2	1	1











































54					1800	2/1	1	1
55		H			3600	4/1	2	2
56		H			1100	8/1	2	1
57		H	PNP		77	4/1	8	2
58		H	OH		1,200	2/1	1	1
59		H	OH		472	1/1	1	1
60					603	2/1	1	1
61		H			3,500	2/1	1	1
62			PNP		44	2/1		
63					40	2/1		
64			PNP	0.56	84	2/2		
65			PNP	0.86	365	1		

























66				0.24	217	1		
67				0.16	20	1		
68				-0.84	52			
69		H			28	4/2	16	2
70		H			93	2/2	1	1
71		H			84	2/1	16	2
72		H	PNP		4	2 <sup>7</sup>		
73		H	OH					
74		H	PNP		34	4/2	1	1
75		H			35	4/4	16	2
76		H			502	4/2	1	1
77		H	PNP		0.4	12/3	24	6




















78		H			3000	1/1	1	1
79		H			3000	2/1	1	1
80		H	OMe		1700	2/1	1	1
81		H		-0.23	53	4/1	1	1
82		H		-0.51	0.1	8/8	32	2
83		H		-1.14	88	4/1	8	2
84		H		0.64	10	2/2	1	1
85		H		-1.11	64	4/1	1	1
86		H		-0.61	826	4/1	1	1
87		H	PNP	-0.51	7	1/1	16	2
88		H	 amide-ester		1,500	1/1	1	1
89		H	 amide		359	4		

90		H	PNP	-0.99	6	1/1	32	4
91		H	PNP	-0.62	5	4/2	32	2
92		H	OH acid	-1.70	446	2		
93		H	PNP	-0.71	9	4/2	4	1
94		H			175	1		
95		H		-0.40	6	1/1	8	1
96		H		-1.06	18	2/1	2	1
97		H	PNP	-0.69	2	8		
98		H		-0.55	39	1/1	2	1
99		H		0.42	3	4/2		
100		H	PNP	-0.31	0.4	5/2		
101		H		-0.91	16	9/1		

























102		H	PNP	-1.23	12	22/2		
103		H	PNP	-1.33	52	4/1		
104		H	PNP	-1.23	5	11/2		
105		H	OH Mono-acid Mono-sodium salt	-0.68	83	8		
106		H		-0.47	0.4	2/1		
107		H		0.38	46	8/1		
108		H		-0.85	234	4/2		
109		H		0.45	9	4/1		
110		H		-2.10	83	4/2		
111		H		0.58	3	4		
112		H		-0.35	60	16		
113		H		-0.24	19	8		
























114		H	PNP	-0.055	0.3	6		
115		H		-0.46	116	16		
116		H		-0.41	48	4		
117		H	PNP	-0.015	0.4	4		
118		H	PNP	-0.31	0.3	4		
119		H	PNP	0.42	1	4		
120			PNP	-1.15	1,000	4		
121			PNP	-0.73	4,000	8		
122		H		-1.08	11	2		
123			PNP	-0.19	101	4		
124			PNP	-0.22	242	4		
125			PNP	-0.065	223	2		























126			PNP	-1.01	244	4		
127		H		0.61	1	1		
128		H	PNP	0.45	0.1	16		
129				-0.85	499	2		
130				-0.90	1,400	4		
131		H		-0.65	29	2		
132		H		0.70	0.4	4		
133		H	PNP	-0.68	6	4		
134		H		-1.08	196	4		
135		H		0.48	6			
136		H		0.095	358	4		
137		H		-0.52	593	4		







138		H			755	2		
139		H	(OBn) <sub>2</sub> Diester			1		
140		H		-0.11	4	2		
141		H		-0.78	655	4		
142		H N-Me	PNP	-0.091	1.3K	2		
143		H N-phenethyl	PNP	1.17	133	4		
144		H N-Et	PNP	-0.036	463	4		
145		H N-benzyl	PNP	0.86	94	4		
146		H N-Me		-0.30	2,500	1		
147		H N-Me		0.84	1,200	1		
148		H		0.13	505	2		
149		H		-0.1	972	1		




150		H N-Me		0.65	765			
151		H		0.23				
152		H		-0.57	96	4		
153		H	PNP	0.17	0.2	16		
154		H N-Me	PNP	0.62	11	4		
155		H	PNP	-0.09	2.0	4		
156		H		-0.28	32	2		
157				-0.51	432	1		
158			PNP	0.13	213	1		
159			PNP	-0.25	71	1		
160				-0.69	219	1		
161				-0.72	140	1		

162				0.28	213	1		
163				-1.30	497			
164		H	PNP	-0.01	0.31	6		
165		H N-Me	PNP	0.71	18	2		
166		H			2,800	1		
167		H	PNP	0.53	2	2		
168		H		0.92	51	1		
169		H		-0.07	140	1		
170		H		0.70?	67	1		
171		H		-0.49	214	1		
172		H		-0.54	1.9K	1		
173		H		-0.14	408	1		

174		H		-0.41	829	1		
175		H		-0.69	9	1		
176		H		-0.31	972	1		
177		H			618	1		
178		H		0.21	185	1		
179		H		-1.10	46			
180		H	PNP	-0.90	1.4			
181		H			1.9K	1		
182		H		-0.98	1.6K	1		
183		H		0.34	205	1		
184		H	PNP	-0.33	0.5	32		
185		H		-0.06	5			

186		H	PNP	0.67	0.2			
187		H			27			
188		H						
189					62	2		

**Table 2:  $\beta$ -Lactamase Inhibition and Microbiological Efficacy of Sulfonamidomethylphosphonate Derivatives.**



Cpd.	R <sup>1</sup>	R <sup>3</sup>	R <sup>4</sup>	Log P	IC <sub>50</sub> ( $\mu$ M) "C" <sup>2</sup>	IC <sub>50</sub> ( $\mu$ M) "A" <sup>3</sup>	IC <sub>50</sub> ( $\mu$ M) "B" <sup>4</sup>	Synergy <sup>5</sup>
								Class C E.CI <sup>6</sup> 10/1 <sup>7</sup>
190		H	PNP	0.58	0.7	>200	>200	8/16
191			OH	0.78	84	>200	>200	0
192			OH	0.58	128	>200	>200	0
193		H	PNP	0.55	4	>200	>200	4

<sup>2</sup> Class C  $\beta$ -lactamase [P99].





















<sup>3</sup> Class A  $\beta$ -lactamase [TEM R+].

















<sup>4</sup> Class B  $\beta$ -lactamase [L-1].



















<sup>5</sup> Expressed as the ratio of the antibiotic MIC determined in the absence of  $\beta$ -lactamase inhibitor to the antibiotic MIC determined in the presence of  $\beta$ -lactamase inhibitor.

<sup>6</sup> E.CI = *Enterobacter cloacae* [ATCC #23355]; H.In. = *Haemophilus influenzae* [ATCC #43163]; St.A. = *Staphylococcus aureus* [MGH:MA #50848].




















<sup>7</sup> The first value provided was determined at an inhibitor concentration of 10  $\mu$ g/mL, and the second value was determined at an inhibitor concentration of 1  $\mu$ g/mL. Where only one value is provided, it was determined at an inhibitor concentration of 10  $\mu$ g/mL.





















194		H		0.37	648	>200	>200	0
195		H	PNP	0.03	0.6	>200	>200	2
196		H	PNP	0.60	0.8	>200	>200	4
197		H		0.64	106	>200		2
198		H		0.66	3	>200	>200	8
199		H		0.72	25	>200	>200	0
200			PNP	0.87	1200	>200	>200	2
201		H			1	>200	>200	8
202		H			594	>200	>200	0
203		H			361	>200	>200	0
204		H			87	>200	>200	0





















205		H			334	>200	>200	8
206		H	PNP	0.85	0.1	467	>200	4
207		H	PNP		1	275	>200	2
208		H	PNP	0.56	3	>200	>200	4
209		H		0.10	142	>200	>200	2
210		H		0.30	23	149	>200	0
211		H			50	>200		0
212		H		1.0	9	64	201	0
213		H			50	14	>200	0
214		H	PNP		0.2	87	>200	16





















215		H			93	494		0
216		H			196	>200	>200	0
217		H			367	>200	>200	
218		H			356	>200	>200	
219		H			619	>200	>200	0
220		H			48	>200	>200	0
221		H	PNP		0.5	>200	>200	8
222		H	PNP		7	>200	>200	4
223		H			9	278	>200	0
224			PNP		183	>200	>200	0









































225		H			1400	>200		0
226		H			0.3	576		16
227		H			9	>200		2
228		H			98	>200		2
229		H	PNP		0.9			
230		H			29			
231		H			165	>200	>200	
232		H			0.2	250	>200	16
233		H			205	>200	>200	0
234		H			219	>200	>200	0





















235		H			5	>200		8
236		H			578	>200	>200	
237		H		0.3	204	>200	>200	0
238		H		-0.4	53	131	>200	0
239		H		0.6	0.5	282	>200	16
240		H		0.1	458	>200		0
241		H			5	>200	>200	2
242		H		-0.2	5	>200	>200	2
243		H		-0.4	6	>200	>200	2
244		H			4	>200	>200	4



















245		H		-0.1	43	272		0
246		H		-0.1	623	>200	>200	2
247		H			0.8	20	>200	8
248		H		0.3	0.06	103		32
249		H			14	>200		2
250		H		0.6	9	>200		2
251		H		-0.6	30	>200		0
252		H			1300	>200		0
253		H		-0.4	0.2	68		16
254		H			0.3	13		32

255		H		0.7	0.6	6		8
256		H			174	>200		0
257		H			354	>200		0
258		H			80	>200		0
259		H			177	>200		0
260		H			130	>200		0
261		H			78	>200		
262		H			165	>252		0
263		H			81	>256		0
264		H			97	394		4























265		H			93	359		4
266		H			82	>244		0
267		H			2	240		2
268		H	PNP	-0.4	3	>240		2
269		H		-0.2	6	107		0
270		H			452	>200		16
271		H	PNP	0.3	2	>200		2
272		H			50	>200		2
273		H			4	54		2
274		H			47	>200		2
275		H			0.1	21		32/8

276		H			63	>200		4
277		H			139	>200		2
278		H			137	>200		2
279		H			109	>200		0
280		H			11	68		0
281		H			4	75		4
282		H			43	662		0
283		H			20	74		0
284		H		-0.3	0.4	511		8

285		H		-0.5	0.4	356		4
286		H			2	>200		16/16
287		H			15	>200		16/8
288		H			375	>200		8/16
289		H			6	>200		32/32
290		H			27	>200		16/32
291		H			666			
292		H			>240	>240		0/0
293		H			1100	>300		0/0
294		H			77	149		4

295		H			27	454		0/0
296		H			98	>200		0/0
297		H			>600	>200		0/0
298		H			0.06	157		32
299		H			2	150		8/8
300		H			28	130		0/2
301		H			4	74		4/4
302		H			3	141		0/2
303		H			9	>200		2/2



304		H			5	>200		16/32
305		H			6	>200		8/4
306		H			7	>200		2/4
307		H			30	>200		2/2
308		H			53	>200		0/0
309					109	>200		0/0
310		H			0.7	19		8/8
311					170	>200		0/0
312		H			0.1	>200		16/8
313		H			27	>200		4/4

***Example 73: Synergistic Effect of  $\beta$ -Lactamase Inhibitors When Tested Against Highly Resistant  $\beta$ -Lactamase Positive Bacterial Strains***







Following procedures identical to those described in Example 72,  $\beta$ -lactamase inhibitors were tested for their ability to enhance antibiotic efficacy against  $\beta$ -lactamase positive bacterial strains that are very highly resistant to  $\beta$ -lactam antibiotics. Representative results are presented in Table 3.

**Highly resistant  $\beta$ -Lactamase positive strains**





Enterobacter cloacae (derepressed)  
Pseudomonas aeruginosa [ATCC#12470-resistant]  
Stenotrophomonas maltophilia [ATCC#12968-resistant]  
Pseudomonas aeruginosa [ATCC#98043010-intermediate resistance]  
Stenotrophomonas maltophilia [ATCC#98043029-intermediate resistance]

Successful inhibition of bacterial  $\beta$ -lactamase activity in these assays is expected to be predictive of success in animals and humans. For examples of the successful clinical development of  $\beta$ -lactamase inhibitors identified by in vitro testing, see, e.g., Di Modugno *et al.*, Current Opinion in Anti-Infective Investigational Drugs 1:26-39 (1999); Moellering, J. Antimicrobial Chemotherapy 31 Suppl. A:1-8 (1993).

Table 3. Synergy results from screening against highly resistant microorganisms.

Cpd.				IC <sub>50</sub> (μM)		Resistant Ent. Cloacae <sup>2</sup>		Resistant Pseudo.		Resistant Stenotro.		Intermed. Pseudo.		Intermed. Stenotro.	
	R <sup>1</sup>	R <sup>3</sup>	R <sup>5</sup>	"C" <sup>1</sup>	10 μg/mL	1 μg/mL	10 μg/mL	1 μg/mL	10 μg/mL	1 μg/mL	10 μg/mL	1 μg/mL	10 μg/mL	1 μg/mL	10 μg/mL
15		H	PNP	18	1		1			1		4		1	
77		H	PNP	0.4	5		1			5		8		1	
72		H	PNP	4	1	1	1	1	1	1	1	1	1	1	1
76		H		502	5		1			5		1		1	

<sup>1</sup> Class C β-lactamase [P99].<sup>2</sup> *Enterobacter Cloacae* (derepressed), *Pseudomonas aeruginosa* (#12470-resistant), *Stenotrophomonas maltophilia* (#12968-resistant), *Pseudomonas aeruginosa* (#98043010-Intermediate resistance), *Stenotrophomonas maltophilia* (#98043029-Intermediate resistance).

82		H		0.1	5		1		5		1		8		1	
114		H	PNP	0.3	5		1		4		1		1		1	
128		H	PNP	0.1	5		1		5		1		2		2	1






























































82		H		0.1	5		1		5		8		1	
114		H	PNP	0.3	5		1		4		1		1	
128		H	PNP	0.1	5	1	1	1	1	1	2	1	2	1

Table 4









































Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
501		H		6	380		2	0	0	0	
502		H		12 15	12K		0	2	0	0	
503		H		30	>250		4	0	0	0	
504		H		1 1	3K		8	2	0	2	
505		H		109 149	126		4	2	0	0	
506		H		37	>250		2	2	0	-	
507		H		11	585		4	2	2	-	
508		H		153 250	>250		2	2	0	0	
509		H		35	69		0	0	0	0	

















Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
510		H		75	140		8	0	0	0	
511		H		45	94		4	0	0	0	
512		H		9	170		4	2	0	0	
513		H		19	>250		4	0	0	0	
514		H		2	259		16	0	0	2	
515		H		8	66		4	2	2	0	
516		H		4 6	628		16	0	0	0	
517		H		9	117		8	0	2	0	
518		H		5 10	227 94		4	0	0	0	
519		H		2 1(MD)	174 98-MD		2	4	4	4	4(oxy)





















Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
520		H		>500	-		4	0	0	0	
521				12 20-MD	407 ≥664		8	2	2	0	
522		H		4 8-MD 1	184 ≥57 26 34		8	2	8	-	2(oxy)
523		H		4 12-MD	62 81		4	2	4	-	
524		H		14	175		2	0	0	-	
525		H		19 6	172		0	2	0	-	
526		H		5 3 (75μM NC) 1-MD 4	1K ≥3K 1K		8	4	8	-	
527		H		55 50 (75μM NC) 183 181	60		4	2	0	-	
528		H		5	162 234		8	2	0	-	



Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
529		H		15 21	128		4	2	2	0	
530		H		56 76	>250		4	2	-	0	
531		H		12	289		4	0	0	0	2oxy
532		H		19 xx-MD	743 ≥52		64	8	2	2	
533		H		7	226		4	4	8	0	
534		H		7	260 557		16	2	2	0	Insol. stock
535		H		28	123		4	4	0	0	
536		H		4	327		4	4	2	2	
537		H		22	150		0	4	0	0	
538		H		2 2 <sub>MD</sub> 1-MD	272 ≥47 >333		64	4	4	0	

Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
539		H		69	272		4	4	0	0	
540		H		20	>250		16	4	4	0	
541		H		1	230		16	4	4	0	
542		H		11 4	37 57		0	4	-	0	
543		H		>250	>250						
544		H					4	0	8	0	
545		H									
546		H		132 125	142 215		2	0	2	0	
547		H		12 8	115 229		16	0	0	0	

Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
548		H		11 10	112 185		4	0	0	0	
549		H		60 34	198 148						
550		H					2	0	0	0	
551		H					2	0	0	0	
552		H		22 <sub>MD</sub>	50		2	0	>16	0	
553		H					2	0	0	0	
554		H					2	0	2	0	
555		H					8	0	0	0	

Cmpd #	R <sub>1</sub>	R	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
556		H		5 <sub>MD</sub>	32		>1 6	0	>16	0	
557		H						0	0		
558		H						0	0		
559		H		6 <sub>MD</sub>	65			0	>16		
560		H									
561		H		61	98						
562		CO <sub>2</sub> Et		>250	>250						
563		CO <sub>2</sub> Et		>250	>250						
719		H									
720		H									


























































Cmpd #	R <sub>1</sub>	R	X	IC <sub>50</sub> (μM) "C"	IC <sub>50</sub> (μM) TEM1 "A"	IC <sub>50</sub> (μM) L-1 "B"	Synergy at 10 μg/ml				LogP or pKa
							Ent. Cl.	Ent. Faec.	Stap. Aureus	P99 Ent. Bact.	
721		H									
722		H									
723		H									
724		H									
725		H									

Table 5



CMPD #	R <sup>1</sup>	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	Synergy (at μg/mL)			
				Class C		Class A	
				E.CI 10/1	Ps. 100	H.In. 10/1	St.A 10
221		F	151	8/8		0/0	
221A.2		F	154	32/4 16/8 32/4	2	0 0 0	0 0
223				4/2			
245		PNP	19				
253				2/0		0	
255				4/2			
276				2/0			
280				2/2		0	0
298				0/0		0/0	
701		PNP	1	8/2		(2)	(2)
702			>100	4/2		(4)	(0)
703			>100	2/2		(8)	(4)

CMPD #	R <sup>1</sup>	X <sup>-</sup>	IC <sub>50</sub> (μM) "C"	Synergy (at μg/mL)			
				Class C		Class A	
				E.CI 10/1	Ps. 100	H.In. 10/1	St.A 10
704		PNP		0/0		0	0
705			255	2/2		(2)	(0)
706			>3K	0/0		(0)	(0)
707			>1K	2/0		(0)	(0)
708		PNP	32	0/0		2	0
709			41	2/0		0	0
710			172	2/0		0	0
711		PNP	207	0/0		0	0
712			>1K	2/0		0	0
713			9	8/2		0	2
714				0/0		0	0
715			2.2K	2/2		0	0

CMPD #	R <sup>1</sup>	X <sup>-</sup>	IC <sub>50</sub> ( $\mu$ M) "C"	Synergy (at $\mu$ g/mL)			
				Class C		Class A	
				E.CI 10/1	Ps. 100	H.In. 10/1	St.A 10
716			316	8/0		0	0
717				8/0		0	0
718		OH	1.6K	4/0		0	0